



ACID-BASE REVIEW KEY

29. Calculate the mass of NaOH which is required to neutralize 25.00 mL of 0.500 M H₂SO₄. (3 marks)




30. A 1.0 M unknown solution was analyzed and the following was observed:



Classify the unknown as an acid or base indicating whether it is weak or strong. Justify your answer using the data provided. (2 marks)

This is a weak acid. It is an acid because bubbles are produced with magnesium. It is weak because the light is dimly glowing.

31. Calculate the pH of 2.0 M nitrous acid. (4 marks)



$$K_a = \frac{x^2}{2.0} = 4.6 \times 10^{-4} \quad 2.0 \rightarrow \sqrt{9.2 \times 10^{-4}} = \sqrt{x^2}$$


$$x = 3.0 \times 10^{-2} \quad \therefore [\text{H}_3\text{O}^+] = 3.0 \times 10^{-2}$$

$$\text{pH} = -\log(3.0 \times 10^{-2})$$

$$\boxed{\text{pH} = 1.52}$$

32. A 2.0 L solution contains one mole of the weak acid, H₃PO₄, in equilibrium with one mole of the salt, NaH₂PO₄. (2 marks)


a) Write an equation that represents this equilibrium.




b) Explain why the pH of this solution does not change significantly when 10.0 mL of 1.014 M KOH is added. (1 mark)

This is a buffer solution, when KOH is added it causes (H₃O⁺) to shift the equilibrium to the products but there is little change in pH!

33. Lactic acid, HC₃H₅O₃, is a compound that accumulates in muscle tissue during exertion. Write the equation and the K_a expression for the ionization of lactic acid in water. (2 marks)






b) Calculate the K_b for HCO_3^- . (1 mark)

$$K_b(\text{HCO}_3^-) = \frac{1.00 \times 10^{-14}}{K_a(\text{H}_2\text{CO}_3)} = \frac{1.00 \times 10^{-14}}{4.3 \times 10^{-7}} = 2.3 \times 10^{-8}$$

37. Calculate the $[\text{H}_3\text{O}^+]$ in 0.550 M $\text{C}_6\text{H}_5\text{COOH}$. (3 marks)



$$K_a = \left(\frac{X^2}{0.550} \right) = 6.5 \times 10^{-5} \Rightarrow 0.550 \rightarrow \sqrt{3.6 \times 10^{-5}} = X^2$$

$$6.0 \times 10^{-3} = X \therefore [\text{H}_3\text{O}^+] = 6.0 \times 10^{-3}$$

34. The ionization constant for water, K_w , is 9.6×10^{-14} at 60°C . (2 marks)



b) Calculate the pH for water at 60°C . (2 marks)

$$K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 9.9 \times 10^{-14} = X^2$$

$$3.09 \times 10^{-7} = X$$

$$\therefore [\text{H}_3\text{O}^+] = 3.09 \times 10^{-7}$$

$$\text{pH} = -\log(3.09 \times 10^{-7})$$

$$\boxed{\text{pH} = 6.51}$$

35. Four monoprotic acids of the same concentration are labelled as follows:

SOLUTION	LABEL
A	$[\text{OH}^-] = 5.0 \times 10^{-11} \text{ M}$
B	$[\text{H}^+] = 0.20 \text{ M}$
C	$\text{pOH} = 11.30$
D	$\text{pH} = 1.20$

List the four solutions in order of decreasing acidity. Use calculations to support your answer. (4 marks)

- (A) $\text{pOH} = -\log(5.0 \times 10^{-11}) = 10.30$; $\text{pH} = 14.000 - 10.30 = 3.70$
- (B) $\text{pH} = -\log(0.20) = 0.70$
- (C) $\text{pOH} = 11.30$; $\text{pH} = 14.000 - 11.30 = 2.70$
- (D) $\text{pH} = 1.20$

$$\boxed{\text{B} > \text{D} > \text{C} > \text{A}}$$


44. What is the $[H_3O^+]$ in a solution formed by adding 60.0 mL of water to 40.0 mL of 0.040 M KOH? (2 marks)

$$[KOH] = \frac{0.040 M \times 0.040 L}{0.100 L} = 0.016 M KOH$$

$$[OH^-] = 0.016 M$$

$$\therefore [H_3O^+] = \frac{1.00 \times 10^{-14}}{0.016 M} = \boxed{6.3 \times 10^{-13}}$$

45. Calculate the pH in 100.0 mL of 0.400 M H_3BO_3 . (4 marks)



assume $0.400 - x \approx 0.400$

$$K_a = \frac{x^2}{0.400} = 1.3 \times 10^{-10} \rightarrow \sqrt{2.92 \times 10^{-10}} = \sqrt{x^2}$$

$$x = 1.7 \times 10^{-5} \therefore [H_3O^+] = 1.7 \times 10^{-5}$$


$$pH = -\log(1.7 \times 10^{-5})$$

$$\boxed{pH = 4.77}$$

46. State two tests that could be safely performed to determine whether an unknown solution is acidic. Predict the results if the solution is acidic. (2 marks)

- Test 1: Test with litmus paper
 Result: should turn Red
- Test 2: Test with magnesium
 Result: should produce bubbles of $H_2(g)$


47. An aqueous solution of Na_2CO_3 undergoes hydrolysis. (1 mark)



b) Calculate K_b for the hydrolysis in a). (1 mark)

$$K_b = \frac{1.00 \times 10^{-14}}{K_a(HCO_3^-)} = \frac{1.00 \times 10^{-14}}{5.6 \times 10^{-11}} = \boxed{1.8 \times 10^{-4}}$$

48. A solution of 0.0100 M lactic acid, $HC_3H_5O_3$, has a pH of 2.95. Calculate the K_a value. (3 marks)



$$pH = 2.95 \therefore [H_3O^+] = \text{antilog}(-2.95) = 1.1 \times 10^{-3}$$

$$K_a = \frac{(1.1 \times 10^{-3})^2}{8.9 \times 10^{-3}} = \boxed{1.4 \times 10^{-4}}$$

49. Calculate the pH of a solution prepared by adding 40.0 mL of 0.440 M NaOH to 60.0 mL of 0.320 M HCl. (3 marks)

$$NaOH + HCl \rightarrow NaCl + H_2O$$

$$[HCl] = \frac{0.320 M \times 0.0600 L}{0.100 L} = 0.192 M H_3O^+$$

$$[NaOH] = \frac{0.440 M \times 0.0400 L}{0.100 L} = 0.176 M OH^-$$

$$[H_3O^+]_{xs} = 0.192 M - 0.176 M$$

$$[H_3O^+]_{xs} = 0.016 M$$

$$pH = -\log(0.016)$$

$$\boxed{pH = 1.80}$$

ACID-BASE REVIEW KEY

38. Calculate the pH of the solution formed by mixing 20.0 mL of 0.500 M HCl with 30.0 mL 0.300 M NaOH. (4 marks)



$$[\text{HCl}]_{\text{ST}} = \frac{0.500 \text{ M} \times 0.0200 \text{ L}}{0.0500 \text{ L}} = 0.200 \text{ M HCl}$$

$$[\text{NaOH}] = \frac{0.300 \text{ M} \times 0.0300 \text{ L}}{0.0500 \text{ L}} = 0.180 \text{ M NaOH}$$

$$[\text{H}_3\text{O}^+]_{\text{XS}} = 0.200 \text{ M} - 0.180 \text{ M} = 0.020 \text{ M}$$

$$\text{pH} = -\log(0.020)$$

$$\boxed{\text{pH} = 1.70}$$

39. a) Write the balanced equation representing the reaction of HF with H₂O. (1 mark)



b) Identify the Brønsted-Lowry bases in the above equation. (1 mark)

H₂O and F⁻

40. Consider the following data:

CHEMICAL SPECIES	FORMULA	IONIZATION CONSTANT
barbituric acid	HC ₄ H ₃ N ₃ O ₃	K _a = 9.8 × 10 ⁻⁵
sodium propionate	NaC ₃ H ₅ O ₂	K _b = 7.5 × 10 ⁻¹⁰
propionic acid	HC ₃ H ₅ O ₂	?


Which is the stronger acid, propionic acid or barbituric acid? Explain, using appropriate calculation. (2 marks)

$$K_a(\text{propionic}) = \frac{1.00 \times 10^{-14}}{7.5 \times 10^{-10}} = 1.3 \times 10^{-5}$$

B/c K_a barbituric acid is > K_a propionic

Barbituric acid is the stronger acid

41. A solution of 0.100 M HOCN has a pH of 2.24. Calculate the K_a value for this acid. (4 marks)



$$\text{pH} = 2.24 \rightarrow [\text{H}_3\text{O}^+] = \text{antilog}(-2.24) = 5.8 \times 10^{-3}$$

$$K_a = \frac{(5.8 \times 10^{-3})^2}{9.4 \times 10^{-2}} = \boxed{3.5 \times 10^{-4}}$$

42. Calculate the pH of a 25.0 mL solution formed by mixing 0.0300 mol HNO₃ and 0.0280 mol NaOH. (2 marks)



$$[\text{H}_3\text{O}^+] = \frac{0.0300 \text{ mol}}{0.0250 \text{ L}} = 1.20 \text{ M}$$

$$[\text{OH}^-] = \frac{0.0280 \text{ mol}}{0.0250 \text{ L}} = 1.12 \text{ M}$$

$$[\text{H}_3\text{O}^+]_{\text{XS}} = 1.20 - 1.12 = 0.08 \text{ M}$$

$$\text{pH} = -\log(0.08)$$

$$\boxed{\text{pH} = 1.1}$$

43. a) Write the net ionic equation for the predominant reaction between NaHSO₃ and NaHCO₃. (2 marks)



b) Explain why the reactants are favoured in the above reaction. (1 mark)

B/c H₂SO₃ is a stronger acid than HCO₃⁻