### DYNAMIC EQUILIBRIUM STUDY GUIDE



**Multiple Choice Section**: This study guide is a compilation of questions from provincial exams since April 1994. I urge you to become intimately familiar with question types. You will notice that questions from one year to another are very similar in their composition. Identification of question types will allow you to be more efficient in answering these questions on the provincial examination. My recommendations for using this study guide are as follows :

- 1. DO ALL THE QUESTIONS in this booklet. These are actual Provincial Exam questions! Your own provincial exam and unit test will include questions similar to the ones in this booklet!
- <u>RESIST THE URGE TO LOOK AT THE ANSWER KEY</u> until you have given all the questions in the section your best effort. Don't do
  one question, then look at the key, then do another and look at the key, and so on. Each time you look at one answer in the study
  guide, your eye will notice other answers around them, and this will reduce the effectiveness of those questions in helping you to
  learn.
- 3. <u>LEARN FROM YOUR MISTAKES</u>! If you get a question wrong, <u>figure out why</u>! If you are having difficulty, <u>talk to your study</u> <u>partner</u>, or maybe <u>phone someone in your Peer Tutoring group</u>. Get together with group members or other students from class and work on these questions together. Explain how you got your answers to tough questions to others. In explaining yourself to someone else, you will learn the material better yourself (try it!) Ask your teacher to explain the questions to you during tutorial or after school. <u>Your goal should be to get 100% on any Chemistry 12 multiple choice test</u>- learning from your mistakes in this booklet will really help you in your efforts to meet this goal!
- 4. This is REALLY CRUCIAL: DO NOT mark the answer anywhere on the questions themselves. For example, do not circle any of options A B C or D-instead use a different sheet of paper to place your answers on. By avoiding this urge, you can re-use this study guide effectively again, when preparing for your final exam. In the box to the left, put an asterisk or small note to yourself to indicate that you got the question wrong and need to come back to it. If you got the question correct initially, a check mark might be assurance that you understand this type of question and therefore can concentrate on other questions that present a challenge to you.
- 5. <u>Check Off the STATUS box on the PRESCRIBED LEARNING OUTCOMES sheet</u>. I have tried to organize the questions in the identical sequence to which they appear on your **Dynamic Equilibrium** Prescribed Learning Outcome sheet. By doing this, you can be confident that you know everything you need to know for both the UNIT EXAM and PROVINCIAL EXAM !

### TABLE OF CONTENTS:

### PART ONE- MULTIPLE CHOICE

INTRODUCTION TO DYNAMIC EQUILIBRIUM	.2
LE CHATELIER'S PRINCIPLE	9
THE EQUILIBRIUM CONSTANT	17
MULTIPLE CHOICE ANSWERS	34
PART TWO- WRITTEN	35
WRITTEN SOLUTION KEY	42

# CHEM....IS....TRY

## INTRODUCTION TO EQUILIBRIUM



**3** D3 Consider the following graph:



# When equilibrium is reached, the <u>rate of the</u> forward reaction is

- A. 0.00 mol/min B. 0.25mol/min
- C. 1.0 mol/min
- D. 3.0 mol/min

4 D3 Consider the following equilibrium:

$$N_{2(g)} + O_{2(g)} \rightleftharpoons 2NO_{(g)}$$

# Nitrogen gas and oxygen gas react when placed in a closed container. As the reaction proceeds towards equilibrium, the <u>rate of the reverse reaction</u>

- A. increases as the concentration of products decreases.
- B. decreases as the concentration of products decreases.
- C. increases as the concentration of products increases.
- D. decreases as the concentration of products increases.
- **5** D3 Consider the following equilibrium:

$$H_2O_{(g)} + CO_{(g)} \rightleftharpoons H_{2(g)} + CO_{2(g)}$$

# At high temperature, $H_2O$ and CO are placed in a closed container. As the system approaches equilibrium, the

A. rate of the forward and reverse reactions both increase.

- B. rate of the forward and reverse reactions both decrease.
- C. rate of the forward reaction decreases and the rate of the reverse reaction increases.
- D. rate of the forward reaction increases and the rate of the reverse reaction decreases.

# 6 D3 A 1.00 L flask contains a gaseous equilibrium system. The addition of reactants to this flask results in a :

- A. shift left and a decrease in the concentration of products.
- B. shift left and an increase in the concentration of products.
- C. shift right and a decrease in the concentration of products.
- D. shift right and an increase in the concentration of products.
- 7 D3 Consider the following:

$$2 \operatorname{NH}_{3(g)} \rightleftharpoons \operatorname{N}_{2(g)} + 3 \operatorname{H}_{2(g)}$$

A flask is initially filled with  $NH_3$ . As the system approaches equilibrium, the rate of the <u>forward</u> reaction

- A. increases as the rate of the reverse reaction decreases.
- B. decreases as the rate of the reverse reaction increases.
- C. increases as the rate of the reverse reaction increases.
- D. decreases as the rate of the reverse reaction decreases.
- 8 D3 Consider the following equilibrium:

$$N_2O_{4(g)} + heat \rightleftharpoons 2NO_{2(g)}$$

Initially, a 1.0 L container is filled with 2.0 mol of  $NO_2$ . As the system approaches equilibrium, the <u>rate of reaction of  $NO_2$ </u>

- A. increases and  $[N_2O_4]$  increases. B. increases and  $[N_2O_4]$  decreases.
- C. decreases and  $[N_2O_4]$  increases. D. decreases and  $[N_2O_4]$  decreases.
- 9 D3 Consider the following equilibrium:

$$\mathrm{SO}_2\mathrm{Cl}_{2(g)} \rightleftharpoons \mathrm{SO}_{2(g)} + \mathrm{Cl}_{2(g)}$$

A 1.0 L container is initially filled with 2.0 mol of  $SO_2CI_2$ . As the reaction proceeds towards equilibrium, the rate of the <u>forward</u> reaction

A. increases and the  $[SO_2]$   $\Box$  increases. B. increases and the  $[SO_2]$  decreases.

- C. decreases and the [SO<sub>2</sub>] increases. D. decreases and the [SO<sub>2</sub>] decreases.
- **10** D3 Consider the following reversible reaction:

$$\operatorname{Fe}_{(aq)}^{3+} + \operatorname{SCN}_{(aq)}^{-} \rightleftharpoons \operatorname{FeSCN}_{(aq)}^{2+}$$

A solution of  $Fe(NO_3)_3$  is added to a solution of KSCN. Which one of the following statements describes the <u>changes</u> in forward and reverse reaction rates <u>as the reaction moves towards</u> <u>equilibrium?</u>

A. Forward and reverse rates increase.

- B. Forward and reverse rates decrease.
- C. Forward rate increases and reverse rate decreases.
- D. Forward rate decreases and reverse rate increases.

**11** D3 Consider the following equilibrium:

$$N_{2(g)} + 2O_{2(g)} \rightleftharpoons 2NO_{2(g)}$$

Equal moles of  $N_2$  and  $O_2$  are added, under certain conditions, to a closed container. Which of the following describes the changes in the <u>reverse</u> reaction which occur as the system <u>proceeds toward equilibrium</u>?

Rate of Reverse Reaction	[NO <sub>2</sub> ]
increases	increases
decreases	increases
increases	decreases
decreases	decreases
	Rate of Reverse Reaction         increases         decreases         increases         decreases         decreases

**12** D3 Consider the following equilibrium:

$$2O_{3(g)} \rightleftharpoons 3O_{2(g)} \qquad K_{eq} = 65$$

Initially 0.10 moles of  $O_3$  and 0.10 moles of  $O_2$  are placed in a 1.0L container. Which of the following describes the <u>changes in concentration</u> as the reaction proceeds towards equilibrium?

	[O <sub>3</sub> ]	[O <sub>2</sub> ]
A.	decreases	decreases
B.	decreases	increases
C.	increases	decreases
D.	increases	increases

**13** D3 Consider the following equilibrium:

 $\mathrm{H_2O}_{(g)} + \mathrm{CO}_{(g)} \ \rightleftarrows \ \mathrm{H_{2(g)}} + \mathrm{CO}_{2(g)}$ 

A closed container is initially filled with  $H_2O$  and CO. As the reaction <u>proceeds towards</u> equilibrium the:

		Α.	[CO] and	I [CO <sub>2</sub> ] both increase. B. [CO] and [CO <sub>2</sub> ] both decrease.		
_		C.	[CO] inc	reases and [CO <sub>2</sub> ] decreases. D. [CO] decreases and [CO <sub>2</sub> ] increases.		
14	D4	W	hich of t	e following describes all chemical equilibrium systems?		
		Α.	The mas	s of the reactants equals the mass of the products.		
		В.	The spe	cies are present in the same ratio as in the balanced equation.		
		C.	The rate	of the forward reaction equals the rate of the reverse reaction.		
		D.	The con	centration of the reactants equals the concentration of the products.		
15	D4	W	nich of t	e following apply to all equilibrium systems?		
			I	Forward and reverse rates are equal		
			II	Macroscopic properties are constant		
			III	Mass of reactants equals mass of products		
_		Α.	I and II c	nly B. I and III only C. II and III only D. I, II and III		

16	D4	Conside	r the following:						
		I I	forward and reverse rates are equal	Which of the above a	re true for <u>all equilibrium</u>				
		II 1	macroscopic properties are constant	<u>systems</u> ?					
		ш	can be achieved from either direction	A. I and II only	B. I and IV only				
		IV G	concentrations of reactants and products are equal	C. I, II and III only	D. II, III and IV only				
17	D4	In <u>all</u> systems at equilibrium, the A. concentration of reactants is less than the concentration of products. B. concentration of reactants and the concentration of products are equal. C. concentration of reactants is greater than the concentration of products.							
18	D4	Consider	r the following equilibrium:		•				
		$2SO_{3(g)} \rightleftharpoons 2SO_{2(g)} + O_{2(g)}$							
	At equilibrium, the rate of decomposition of SO <sub>3</sub>								
		A. equals	s the rate of formation of $O_2$ B. 6 s than the rate of formation of $O_2$ D i	equals the rate of formation	$1 \text{ of } SO_3$				
19	D4	Which o	f the following statements are true for	all equilibrium systems					
		I. Macros	scopic properties are constant.	_ · _ /					
		II. Mass	of the reactants equals mass of the prod	ucts.					
		III. An ec	quilibrium can be achieved from either pro	oducts or reactants.					
		A. I and	II only B. I and III only C. II and III or	nly D. I, II and III					
20	D4	Which of the following is true for <u>all</u> equilibrium systems?							
		A. The m	hass of reactants is equal to the mass of	products.					
		C The c	oncentration of reactants is equal to the	concentration of products					
		D. The ra	ate of the forward reaction is equal to the	rate of the reverse reactio	n.				
21	D4	Which o	f the following is characteristic of <u>all</u> s	systems at equilibrium?					
		A. Activation energy is not required.							
		B. Changes do not occur at the microscopic level.							
		D. Temperature and pressure affect the equilibrium position equally.							
22	D4	Conside	er the following:						
		[							
		Ι	constant temperature						
		II	equal concentrations of reactants a	nd products					

III equal rates of forward and reverse reactions

### A system at equilibrium <u>must</u> have

		A. I and II only	B. I and III only	C. II and III only	D. I, II and III.
23	D4	Macroscopic proper	ties become <u>constant</u> in a	n equilibrium system <u>wh</u>	en
		A. all reactions have s	stopped.	B. the reactants are c	ompletely used up.
		C. maximum enthalpy	has been reached.	D. forward and revers	e reaction rates are equal.
24	D4	Which of the followi	ng does <u>not a</u> pply to all cl	nemical equilibrium syste	ms? AUG 2000
		A. They are closed			
		B. The macroscopic p	properties are constant.		
		C. Forward and rever	se reaction rates are equal.		
		D There are equal co	incentrations of reactants ar	nd products	

25	D4	Which	of the following applies	to a <u>chemical equilib</u>	rium?			
		I.	Forward and reverse react	ion rates are equal	]			
		II.	Equilibrium can be achiev	ed from either direction				
		III.	Macroscopic properties ar	e constant				
_		A. I onl	y B. I and II only	C. II and III only	D.	I, II and III		
26	D5	Chemic A. the r C. the r	cal equilibrium is said to eaction proceeds quickly. nacroscopic properties ar	b be <u>dynamic</u> because B. the r re constant. D. both	) nass o forwai	f the reactants is or rd and reverse rea	decreasing. Ictions are occurrir	ng.
27	D5	<ul> <li>Equilibrium is a <u>dynamic</u> process because the</li> <li>A. macroscopic properties are not changing.</li> <li>B. mass of reactants equals the mass of products.</li> <li>C. forward and reverse reactions continue to occur.</li> </ul>						
28	D5	Equilib	prium is said to be dyna	mic because the	stant.			
		A. forward A. forward C. forward	ard and reverse reactions ard reaction goes to com	stop B. reve pletion D. forw	rse rea ard an	action goes to com d reverse reaction	pletion. s continue.	
29	D5	<ul> <li>A system at equilibrium is said to be <u>dynamic</u> because at equilibrium the</li> <li>A. temperature does not change.</li> <li>B. macroscopic properties are constant.</li> <li>C. forward and reverse reactions continue to occur.</li> <li>D. concentrations of reactants and products are constant.</li> </ul>						
30	D5	<ul> <li>A chemical equilibrium is described as <u>dynamic</u> because :</li> <li>A. maximum randomness has been achieved.</li> <li>B. the pressure and temperature do not change.</li> <li>C. both reactants and products continue to form.</li> <li>D. the concentrations of chemical species remain constant.</li> </ul>						
31	D7	Consid	er the following reaction:					1
		$N_{2}$	$(g) + 3H_{2(g)} \rightarrow 2NH_{3(g)}$	(g) + energy		ENTHALPY	ENTROPY	
					Δ	increases	decreases	

Which of the following describes the changes in <u>enthalpy</u> and <u>entropy</u> as the <u>reaction proceeds</u>?

	ENTHALPY	ENTROPY
А.	increases	decreases
В.	increases	increases
C.	decreases	decreases
D.	decreases	increases

- **32** D7 In which reaction is entropy decreasing?
  - A.  $H_2O_{(\ell)} \rightarrow H_2O_{(g)}$
  - B.  $N_2O_{4(g)} \rightarrow 2NO_{2(g)}$

C. 
$$\operatorname{CaCO}_{3(s)} \to \operatorname{CaO}_{(s)} + \operatorname{CO}_{2(g)}$$

D. 
$$\operatorname{Fe}^{3+}(aq) + \operatorname{SCN}^{-}(aq) \to \operatorname{FeSCN}^{2+}(aq)$$

**33** D7 Consider the following equilibrium:

$$N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + 92 \text{ kJ}$$

The forward reaction is

A. exothermic and entropy is increasing.	B. exothermic and entropy is decreasing.
C. endothermic and entropy is increasing.	D. endothermic and entropy is decreasing.

### 34 D7 In which reaction is the enthalpy of the reactants greater than the enthalpy of the products?

- A.  $H_2O_{(s)} \rightarrow H_2O_{(\ell)}$ B.  $H_2O_{(s)} \rightarrow H_2O_{(g)}$
- **D.**  $\Pi_2 O(s) \to \Pi_2 O(g)$
- C.  $H_2O_{(\ell)} \rightarrow H_2O_{(s)}$
- D.  $H_2O_{(\ell)} \rightarrow H_2O_{(g)}$
- 35 D7 Consider the following reaction:

$$Na_{2}CO_{3(s)} + 2HCl_{(aq)} \rightarrow 2NaCl_{(aq)} + CO_{2(g)} + H_{2}O_{(\ell)} \qquad \Delta H = -27.7 \text{ kJ}$$

### In this reaction,

- A. minimum enthalpy and maximum entropy both favour products.
- B. minimum enthalpy and maximum entropy both favour reactants.
- C. minimum enthalpy favours products and maximum entropy favours reactants.
- D. minimum enthalpy favours reactants and maximum entropy favours products.
- **36** D7 Consider the following possible reaction:

$$N_2O_{(g)} + NO_{2(g)} \rightarrow 3NO_{(g)} \qquad \Delta H = +156 \text{ kJ}$$

### Which of the following statements is correct?

- A. Minimum enthalpy and maximum entropy both favour the products.
- B. Minimum enthalpy and maximum entropy both favour the reactants.
- C. Minimum enthalpy favours the reactants and maximum entropy favours the products.
- D. Minimum enthalpy favours the products and maximum entropy favours the reactants.
- 37 D7 Consider the following equilibrium:

$$2NO_{2(g)} \rightleftharpoons N_2O_{4(g)} + 59 \text{ kJ}$$

For the above reaction,

A. both minimum enthalpy and maximum entropy favour products.

- B. both minimum enthalpy and maximum entropy favour reactants.
- C. minimum enthalpy favours reactants and maximum entropy favours products.
- D. minimum enthalpy favours products and maximum entropy favours reactants.
- 38 D7 In which of the following does the entropy decrease?

A. 
$$\operatorname{NaCl}_{(s)} \to \operatorname{Na}_{(aq)}^+ + \operatorname{Cl}_{(aq)}^-$$

B. 
$$4NO_{(g)} + 6H_2O_{(g)} \rightarrow 4NH_{3(g)} + 5O_{2(g)}$$

C. 
$$2 \operatorname{NaHCO}_{3(s)} \rightarrow \operatorname{Na_2CO}_{3(s)} + \operatorname{CO}_{2(g)} + \operatorname{H_2O}_{(g)}$$

D. 
$$\operatorname{CaCO}_{3(s)} + 2\operatorname{HCl}_{(aq)} \rightarrow \operatorname{CaCl}_{2(aq)} + \operatorname{CO}_{2(g)} + \operatorname{H}_2\operatorname{O}_{(\ell)}$$

# 39 D7 In which of the following systems will the factors of entropy and enthalpy <u>both</u> favour the <u>reactants</u>?

- A.  $3C_{(s)} + 3H_{2(g)} + heat \rightleftharpoons C_3H_{6(g)}$
- B.  $PCl_{5(g)} + heat \rightleftharpoons PCl_{3(g)} + Cl_{2(g)}$
- C.  $\text{NH}_4\text{Cl}_{(s)} + \text{heat} \rightleftharpoons \text{NH}_{4(aq)}^+ + \text{Cl}_{(aq)}^-$
- D.  $\operatorname{Cl}_{2(g)} + 2\operatorname{HI}_{(g)} \rightleftharpoons \operatorname{I}_{2(g)} + 2\operatorname{HCl}_{(g)} + \operatorname{heat}$

40	D7	Consider the following reaction:			AUG 2000
		$C_3H_{8(g)} + 5O_{2(g)} \rightarrow 3CO_{2(g)} + 4H_2O_{(g)}$	$\Delta H =$	-2202 kJ	
				Entropy	Enthalpy
		Which of the following applies to the	А.	increases	increases
		forward reaction?	B.	increases	decreases
			C.	decreases	increases
			D.	decreases	decreases

41 D7 Which of the following reactions results in an <u>entropy increase</u>?

- - -

A. 
$$2C_{(s)} + O_{2(g)} \rightarrow 2CO_{(g)}$$
  
B.  $N_{2(g)} + 2H_{2(g)} \rightarrow N_2H_{4(\ell)}$   
C.  $2SO_{2(g)} + O_{2(g)} \rightarrow 2SO_{3(g)}$   
D.  $Ag^+_{(aq)} + Cl^-_{(aq)} \rightarrow AgCl_{(s)}$ 

#### 42 D8 In an endothermic equilibrium system, the

- A. minimum enthalpy and the maximum entropy both favour products.
- B. minimum enthalpy and the maximum entropy both favour reactants.
- C. minimum enthalpy favours products and the maximum entropy favours reactants.
- D. minimum enthalpy favours reactants and the maximum entropy favours products.

#### 43 D8 Chemical systems tend to move toward positions of

- A. minimum enthalpy and maximum entropy.
- B. maximum enthalpy and minimum entropy.
- C. minimum enthalpy and minimum entropy.
- D. maximum enthalpy and maximum entropy.
- **44** D9 Consider the enthalpy and entropy changes in the following:

$$C_2H_{2(g)} + H_{2(g)} \xrightarrow{?} C_2H_{4(g)} \qquad \Delta H = -175 \text{ kJ}$$

#### Which of the following statements is correct?

- A. No reaction occurs because both the enthalpy and entropy factors favour the reactants.
- B. The reaction goes to completion because both the enthalpy and entropy factors favour the product.
- C. The system reaches equilibrium because the enthalpy factor favours the reactants and the entropy factor favours the product.
- D. The system reaches equilibrium because the enthalpy factor favours the product and the entropy factor favours the reactants.

# 45 D9 In which of the following reactions does the tendency towards minimum enthalpy and maximum entropy <u>oppose</u> each other?

А.	$3O_{2(g)} \rightarrow 2O_{3(g)}$	$\Delta H = +285 \text{ kJ}$
B.	$\tfrac{1}{2}\operatorname{N}_{2(g)} + \operatorname{O}_{2(g)} \to \operatorname{NO}_{2(g)}$	$\Delta H = +34 \text{ kJ}$
C.	$2\mathrm{H}_2\mathrm{O}_{(g)} \to 2\mathrm{H}_{2(g)} + \mathrm{O}_{2(g)}$	$\Delta H = +484 \text{ kJ}$
D.	$\mathrm{P}_{4(s)} + 6\mathrm{H}_{2(g)} \to 4\mathrm{PH}_{3(g)}$	$\Delta H = +37 \text{ kJ}$

46	D9	in wh maxii A.	from the following systems would the tendencies mum entropy be in <u>opposition</u> to each other? $Br_{2(l)} + heat \rightarrow Br_{2(g)}$	s toward minimum enthalpy and
		В.	$\operatorname{NaOH}_{(s)} \to \operatorname{Na}_{(aq)}^+ + \operatorname{OH}_{(aq)}^- + \operatorname{heat}$	
		C. 2	$2C_{(g)} + 2H_{2(g)} \rightarrow C_2H_{4(g)}$ $\Delta H$ is positive	
		D.	$K_{(s)} + H_2O_{(1)} \to K^+_{(aq)} + OH^{(aq)} + \frac{1}{2}H_{2(g)}$ $\Delta H$ is	s negative
47	D9	In wh react	ich of the following do both minimum enthalpy ar <u>ants</u> ?	nd maximum entropy factors <u>favor the</u>
		Α.	$\operatorname{Cl}_{2(g)} \rightleftharpoons \operatorname{Cl}_{2(aq)}$	$\Delta H = -25  kJ$
		В.	$C_{(s)} + H_2O_{(\ell)} \rightleftharpoons CO_{(g)} + H_{2(g)}$	$\Delta H = +131  kJ$
		C.	$2\mathrm{CO}_{2(g)} + 3\mathrm{H}_2\mathrm{O}_{(g)} \rightleftharpoons \mathrm{C}_2\mathrm{H}_5\mathrm{OH}_{(\ell)} + 3\mathrm{O}_{2(g)}$	$\Delta H = +1239  kJ$
		D.	$Na_2CO_{3(s)} + HCl_{(aq)} \rightleftharpoons 2NaCl_{(aq)} + CO_{2(g)} + H_2$	$_{2}O_{(\ell)}$ $\Delta H = -28  \text{kJ}$
48	D9	In wh oppo	ich of the following will the driving forces of mini <u>se</u> one another?	mum enthalpy and maximum entropy
		A.	$2C_{(s)} + O_{2(g)} \rightarrow 2CO_{(g)}$	$\Delta H = -221  kJ$
		B.	$2\mathrm{N}_{2(g)} + \mathrm{O}_{2(g)} \rightarrow 2\mathrm{N}_{2}\mathrm{O}_{(g)}$	$\Delta H = +164 \text{ kJ}$
		C.	$2\mathrm{CO}_{(g)} + \mathrm{O}_{2(g)} \to 2\mathrm{CO}_{2(g)}$	$\Delta H = -566  kJ$
		D.	$4\mathrm{CO}_{2(g)} + 6\mathrm{H}_2\mathrm{O}_{(g)} \rightarrow 2\mathrm{C}_2\mathrm{H}_{6(g)} + 7\mathrm{O}_{2(g)}$	$\Delta H = +3122 \text{ kJ}$

### LE CHATELIER'S PRINCIPLE

Consider the following equilibrium: 49 E2  $4 \text{NH}_{3(g)} + 5 \text{O}_{2(g)} \rightleftharpoons 4 \text{NO}_{(g)} + 6 \text{H}_2 \text{O}_{(g)} + \text{energy}$ Which of the following will cause the equilibrium to shift to the left? A. adding  $H_2O_{(g)}$   $\square$   $\square$   $\square$  B. removing some  $NO_{(g)}$ C. increasing the volume D. decreasing the temperature Consider the following equilibrium: 50 E2  $2NO_{(g)} + O_{2(g)} \rightleftharpoons 2NO_{2(g)} + energy$ When the volume of the container is increased, the equilibrium shifts to the A. left and K <sub>eq</sub> decreases. B. right and  $K_{eq}$  increases. C. left and K<sub>eq</sub> remains constant. D. right and K eq remains constant. Consider the following equilibrium: 51 E2  $2SO_{2(g)} + O_{2(g)} \rightleftharpoons 2SO_{3(g)} + energy$ 

Which of the following will cause this equilibrium to shift to the left?A. adding a catalystB. adding some SO2C. increasing the volumeD. decreasing the temperature

**52** E2 Consider the following equilibrium:

$$2NO_{(g)} + Br_{2(g)} + energy \rightleftharpoons 2NOBr_{(g)}$$

### The equilibrium will shift to the left as a result of:

A. adding a catalyst. B. removing NOBr. C. increasing the volume. D. increasing the temperature. 53 E2 Consider the following equilibrium:

$$N_{2(g)} + O_{2(g)} + energy \rightleftharpoons 2NO_{(g)}$$

#### When the temperature is increased, the equilibrium shifts to the

A. left and K <sub>eq</sub> increases.	B. left and K <sub>eq</sub> decreases.
C. right and $K_{eq}$ increases.	D. right and K <sub>eq</sub> decreases.

54 E2 Consider the following equilibrium:

$$2 \operatorname{NO}_{2(g)} \rightleftharpoons \operatorname{N}_2 \operatorname{O}_{4(g)} + \operatorname{energy}$$

### The equilibrium will shift to the left as a result of

A. adding a catalyst.B. increasing the volume.C. removing some  $N_2O_4$ .D. decreasing the temperature.

55 E2 Consider the following equilibrium:

$$C_{(s)} + 2H_{2(g)} \rightleftharpoons CH_{4(g)}$$

### <u>The addition of $H_2$ will cause the equilibrium to shift to the</u>

A. left and [CH <sub>4</sub> ] will increase.	B. left and [CH₄] will decrease.
C. right and [CH <sub>4</sub> ] will increase.	D. right and [CH <sub>4</sub> ] will decrease.
Consider the fellowing equilibrium.	

**56** E2 Consider the following equilibrium:

$$\operatorname{PCl}_{5(g)} \rightleftharpoons \operatorname{PCl}_{3(g)} + \operatorname{Cl}_{2(g)}$$

The equilibrium concentration of PCI<sub>5</sub> will increase when

A. PCl<sub>3</sub> is addedB. Cl<sub>2</sub> is removedC. a catalyst is addedD. the volume of the container is increased.Consider the following equilibrium:

$$N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)}$$

### If the volume of the container is decreased, the

A.  $K_{eq}$  decreases B.  $[N_2O_4]$  increases C. equilibrium does not shift D. equilibrium shifts to the right. 58 E2 Consider the following equilibrium:

$$\mathrm{NH}_{3(g)} + \mathrm{HCl}_{(g)} \rightleftharpoons \mathrm{NH}_4\mathrm{Cl}_{(s)} + \mathrm{energy}$$

### Which of the following will result in a decrease in the mass of NH<sub>4</sub>CI ?

A. adding NH<sub>3</sub> B. removing HCl C. decreasing the volume D. decreasing the temperature

57

E2

**59** E2 Consider the following equilibrium:

$$H_{2(g)} + I_{2(g)} \rightleftharpoons 2HI_{(g)}$$

# The pressure on the system is increased by reducing the volume. When <u>comparing the</u> <u>new equilibrium with the original equilibrium</u>,

A. all concentrations remain constant.

- B. the concentrations of all species have increased.
- C. reactant concentrations have increased while product concentrations have decreased.
- D. reactant concentrations have decreased while product concentrations have increased.

60 E2 Consider the rate diagram below for the following reaction:



$$2 \operatorname{HI}_{(g)} \rightleftharpoons \operatorname{H}_{2(g)} + \operatorname{I}_{2(g)}$$

### Which of the following occurs at time $t_1$ ?

- A. addition of  $H_2$
- B. addition of HI

C. addition of a catalyst

D. a decrease in volume

**61** E2 Given the following system:

$$2 \operatorname{CrO}_{4}^{2-}_{(aq)} + 2 \operatorname{H}_{(aq)}^{+} \rightleftharpoons \operatorname{Cr}_{2} \operatorname{O}_{7}^{2-}_{(aq)} + \operatorname{H}_{2} \operatorname{O}_{(\ell)}$$

Which of the following chemicals, when added to the above system at equilibrium, would result in a <u>decrease</u> in  $[CrO_4^2]$ ?

A. NaOHB.  $HNO_3$ C.  $Na_2CrO_4$ D.  $Na_2Cr_2O_7$ E2Consider the following equilibrium:

$$SO_{2(g)} + NO_{2(g)} \rightleftharpoons SO_{3(g)} + NO_{(g)} + energy$$

The equilibrium does <u>not</u> shift with a change in the

A. volume.B. temperature.C. concentration of products.D. concentration of reactants.E2Consider the following equilibrium:

$$SO_2Cl_{2(g)} + energy \rightleftharpoons SO_{2(g)} + Cl_{2(g)}$$

When the temperature is decreased, the equilibrium shifts

	<u></u> ,	
A. left and [SO <sub>2</sub> Cl <sub>2</sub> ] increases.	B. left and [SO <sub>2</sub> Cl <sub>2</sub> ] decreases.	
C. right and [SO <sub>2</sub> Cl <sub>2</sub> ] increases.	D. right and [SO <sub>2</sub> Cl <sub>2</sub> ] decreases.	
Consider the following equilibrium		

$$N_2O_{4(g)} + 58 \text{ kJ} \rightleftharpoons 2NO_{2(g)}$$

The equilibrium shifts right when

A. NO <sub>2</sub> is added	B. N <sub>2</sub> O <sub>4</sub> is removed
C. the temperature is decreased	D. the volume of the system is increased.

62

63

64

E2

65	E2	Consider the following equilibrium:
		$CH_{4(g)} + H_2O_{(g)} + heat \rightleftharpoons CO_{(g)} + 3H_{2(g)}$
		In which of the following will <u>both stresses</u> shift the <u>equilibrium right</u> ?
		A. a decrease in temperature and a decrease in volume
		B. an increase in temperature and a decrease in volume
		D. an increase in temperature and an increase in volume
66	E2	Consider the following equilibrium:
		$2 \operatorname{HI}_{(g)} \rightleftharpoons \operatorname{H}_{2(g)} + \operatorname{I}_{2(g)} \qquad \Delta \operatorname{H} = -68 \text{ kJ}$
		Which of the following would <u>cause</u> the <u>equilibrium to shift right</u> ?
		A. Increasing the volume. B. Decreasing the volume.
67	E2	Consider the following equilibrium:
01		
		$2SO_{2(g)} + O_{2(g)} \rightleftharpoons 2SO_{3(g)}$
		Which of the following will <u>shift</u> the <u>equilibrium to the right</u> ?
		I. adding more O 2
		III. adding a catalyst
68	F2	A. I only B. III only C. I and II only D. II and III only
00	66	
		$2 \operatorname{HI}_{(g)} \rightleftharpoons \operatorname{H}_{2(g)} + \operatorname{I}_{2(g)}$
		At constant temperature and volume, more $I_2$ is added to the above equilibrium. A <u>new</u> state of
		equilibrium results from a shift to the A left with a net decrease in [Ha] B left with a net increase in [Ha]
		C. right with a net increase in $[H_2]$ . D. right with a net decrease in $[H_2]$ .
69	E2	When the temperature of an equilibrium system is increased, the equilibrium always shifts to
		favour the
70	E2	A exothermic reaction. D. endothermic reaction. C. formation of products. D. formation of reactants.
		Consider the following equilibrium:
		Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + 92 \text{ kJ}$
		Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + 92 \text{ kJ}$ In which of the following will both changes <u>shift the equilibrium right</u> ?
		Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + 92 \text{ kJ}$ In which of the following will both changes <u>shift the equilibrium right</u> ? A. An increase in volume and a decrease in temperature. B. An increase in volume and an increase in temperature.
		Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + 92 \text{ kJ}$ In which of the following will both changes <u>shift the equilibrium right</u> ? A. An increase in volume and a decrease in temperature. B. An increase in volume and an increase in temperature. C. A decrease in volume and a decrease in temperature.
		Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + 92 \text{ kJ}$ In which of the following will both changes <u>shift the equilibrium right</u> ? A. An increase in volume and a decrease in temperature. B. An increase in volume and an increase in temperature. C. A decrease in volume and a decrease in temperature. D. A decrease in volume and an increase in temperature.
71	E2	Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + 92 \text{ kJ}$ <b>In which of the following will both changes <u>shift the equilibrium right</u>? A. An increase in volume and a decrease in temperature. B. An increase in volume and an increase in temperature. C. A decrease in volume and a decrease in temperature. D. A decrease in volume and an increase in temperature. Consider the following equilibrium:</b>
71	E2	Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + 92 \text{ kJ}$ In which of the following will both changes <u>shift the equilibrium right</u> ? A. An increase in volume and a decrease in temperature. B. An increase in volume and an increase in temperature. C. A decrease in volume and a decrease in temperature. D. A decrease in volume and an increase in temperature. Consider the following equilibrium: $H_{2(g)} + CO_{2(g)} \rightleftharpoons CO_{(g)} + H_2O_{(g)} \qquad \Delta H = +41 \text{ kJ}$
71	E2	Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + 92 \text{ kJ}$ <b>In which of the following will both changes <u>shift the equilibrium right</u>? A. An increase in volume and a decrease in temperature. B. An increase in volume and an increase in temperature. C. A decrease in volume and a decrease in temperature. D. A decrease in volume and an increase in temperature. Consider the following equilibrium: <math>H_{2(g)} + CO_{2(g)} \rightleftharpoons CO_{(g)} + H_2O_{(g)} \qquad \Delta H = +41 \text{ kJ}</math> The temperature of the above equilibrium system is <u>increased</u> while kept at a constant volume.</b>
71	E2	Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + 92 \text{ kJ}$ In which of the following will both changes <u>shift the equilibrium right</u> ? A. An increase in volume and a decrease in temperature. B. An increase in volume and an increase in temperature. C. A decrease in volume and a decrease in temperature. D. A decrease in volume and an increase in temperature. Consider the following equilibrium: $H_{2(g)} + CO_{2(g)} \rightleftharpoons CO_{(g)} + H_2O_{(g)} \qquad \Delta H = +41 \text{ kJ}$ The temperature of the above equilibrium system is <u>increased</u> while kept at a constant volume. A new state of equilibrium is established in which there is
71	E2	Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + 92 \text{ kJ}$ In which of the following will both changes <u>shift the equilibrium right</u> ? A. An increase in volume and a decrease in temperature. B. An increase in volume and an increase in temperature. C. A decrease in volume and a decrease in temperature. D. A decrease in volume and an increase in temperature. Consider the following equilibrium: $H_{2(g)} + CO_{2(g)} \rightleftharpoons CO_{(g)} + H_2O_{(g)} \qquad \Delta H = +41 \text{ kJ}$ The temperature of the above equilibrium system is <u>increased</u> while kept at a constant volume. A new state of equilibrium is established in which there is A. an increase in [CO] and a decrease in $K_{eq}$ B. an increase in [CO] and an increase in $K_{eq}$ C. an increase in [CO] and a decrease in $K_{eq}$ D. an increase in [CO] and an increase in $K_{eq}$
71		Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + 92 \text{ kJ}$ In which of the following will both changes <u>shift the equilibrium right</u> ? A. An increase in volume and a decrease in temperature. B. An increase in volume and an increase in temperature. C. A decrease in volume and a decrease in temperature. D. A decrease in volume and an increase in temperature. Consider the following equilibrium: $H_{2(g)} + CO_{2(g)} \rightleftharpoons CO_{(g)} + H_2O_{(g)} \qquad \Delta H = +41 \text{ kJ}$ The temperature of the above equilibrium system is <u>increased</u> while kept at a constant volume. A new state of equilibrium is established in which there is A. an increase in [CO] and a decrease in K <sub>eq</sub> B. an increase in [CO] and an increase in K <sub>eq</sub>
71	E2	Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + 92 \text{ kJ}$ In which of the following will both changes <u>shift the equilibrium right</u> ? A. An increase in volume and a decrease in temperature. B. An increase in volume and an increase in temperature. C. A decrease in volume and a decrease in temperature. D. A decrease in volume and an increase in temperature. Consider the following equilibrium: $H_{2(g)} + CO_{2(g)} \rightleftharpoons CO_{(g)} + H_2O_{(g)}$ $\Delta H = +41 \text{ kJ}$ The temperature of the above equilibrium system is <u>increased</u> while kept at a constant volume. A new state of equilibrium is established in which there is A. an increase in [CO] and a decrease in $K_{eq}$ B. an increase in [CO] and an increase in $K_{eq}$ C. an increase in [CO <sub>2</sub> ] and a decrease in $K_{eq}$ D. an increase in [CO <sub>2</sub> ] and an increase in $K_{eq}$

72 E2 Consider the following equilibrium:

$$2SO_{3(g)} \rightleftharpoons 2SO_{2(g)} + O_{2(g)}$$

The <u>volume</u> of the system is <u>decreased</u> at a <u>constant temperature</u>. A new state of equilibrium is established by a shift of the <u>original</u> equilibrium to the

A. left and [SO<sub>3</sub>] increases. B. right and [SO<sub>3</sub>] decreases.

C. left and [SO<sub>3</sub>] remains unchanged. D. right and [SO<sub>3</sub>] remains unchanged.

**73** E2 Consider the following equilibrium system:

$$\operatorname{NH}_{3(aq)} + \operatorname{H}_2\operatorname{O}_{(l)} \rightleftharpoons \operatorname{NH}_{4(aq)}^+ + \operatorname{OH}_{(aq)}^-$$

Which of the following <u>when added</u> to the above equilibrium system, would cause an <u>increase</u> in [OH] ?

**A**.  $NH_3$  **B**.  $H_2O$  **C**.  $NH_4^+$  **D**. HCl **74** E2 Consider the following graph which relates to this equilibrium:

$$N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} \qquad \Delta H = -92 \text{ kJ}$$

Which of the following <u>caused</u> the changes in the concentrations at time *t* ?

- A. addition of N<sub>2</sub>
- B. removal of  $H_2$
- C. decrease in temperature
- D. decrease in reaction volume



**75** E2 Consider the following equilibrium:

$$C_{(s)} + 2H_{2(g)} \rightleftharpoons CH_{4(g)} + 74 \text{ kJ}$$

When a small amount of solid C is added to the system,

A. [H2] decreasesB. [CH4] increases.C. the temperature increasesD. all concentrations remain constant.

**76** E2 Consider the following equilibrium system:

$$\operatorname{CO}_{2(g)} + \operatorname{H}_{2(g)} \rightleftharpoons \operatorname{CO}_{(g)} + \operatorname{H}_2\operatorname{O}_{(g)}$$

Which of the following when added to the system above would result in a net decrease in [H<sub>2</sub>O] A.  $CO_2$  B.  $H_2$  C. CO D.  $H_2O$ 

E2 Which of the following reactions will shift left when pressure is increased <u>and</u> when temperature is decreased?

A. 
$$N_{2(g)} + 2O_{2(g)} + heat \rightleftharpoons 2NO_{2(g)}$$

B. 
$$N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + heat$$

C. 
$$CH_{4(g)} + H_2O_{(g)} + heat \rightleftharpoons CO_{(g)} + 3H_{2(g)}$$

D. 
$$CS_{2(g)} + 4H_{2(g)} \rightleftharpoons CH_{4(g)} + 2H_2S_{(g)} + heat$$

78 E2 Consider the following equilibrium system:

$$\operatorname{FeO}_{(s)} + \operatorname{H}_{2(g)} \rightleftharpoons \operatorname{Fe}_{(s)} + \operatorname{H}_{2}\operatorname{O}_{(g)}$$

Which one of the following statements describes the effect that a <u>decrease in volume</u> would have on the position of equilibrium <u>and</u> the [H<sub>2</sub>] in the above system?

A. No shift, [H <sub>2</sub> ] increases.	B. Shift right, [H <sub>2</sub> ] increases.
C. Shift right, [H <sub>2</sub> ] decreases.	D. No shift, [H <sub>2</sub> ] remains constant.

77

79	E2	Consider the following equilibrium system:	
		$CaCO_{3(s)} \rightleftharpoons CaO_{(s)} + CO_{2(g)}$	
		<b>Which one of the following <u>changes</u> would cause the above system to <u>shift left</u>? A. Add more CaO . B. Remove CaCO<sub>3</sub> C. Decrease volume D. Increase surface area of CaO .</b>	
80	E2	Consider the following concentration versus time graph for the equilibrium:	
		$N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)}$	
		At time = " t ", which one of the following <u>stresses</u> occurred?	
		A. Catalyst was added. B. Pressure was changed. C. Temperature was changed. D. Concentration of NO <sub>2</sub> was changed.	
		"t" TIME(s)	
81	E2	Consider the following equilibrium:	
		$H_{2(g)} + I_{2(g)} \rightleftharpoons 2HI_{(g)} \qquad [HI] \qquad [HI] \qquad [HI]$	
		Which graph represents what happens time time time	
		C. [HI] D. [HI] [HI] [HI] [HI] [HI]	
82	E2	Consider the following equilibrium:	
		$CO_{(g)} + H_2O_{(g)} \rightleftharpoons CO_{2(g)} + H_{2(g)} \qquad \Delta H = -41  \text{kJ}$	
		What will cause a shift in the equilibrium?           A. adding a catalyst         B. changing volume         C. adding an inert gas         D. changing temperature	
83	ΕZ	Consider the following equilibrium: $CH_3COOH_{(aq)} + H_2O_{(\ell)} \rightleftharpoons CH_3COO^{(aq)} + H_3O^+_{(aq)} + heat$	
		A stress was applied at time $t_1$ and the data was plotted on the following graph:	
		The stress that was imposed at time $t_1$ is the result of: A. the addition of HCl.	
		B. decreasing the temperature.	
		D. Interaction of NaCh COU 3.	

D. increasing the volume of the container.

84 E2 Consider the following equilibrium:

$$2CO_{(g)} + O_{2(g)} \rightleftharpoons 2CO_{2(g)} + energy$$

Some  $CO_2$  is added to the equilibrium system at constant volume and a new equilibrium is established. Compared to the original equilibrium, the <u>rates of the forward and reverse reactions for</u> <u>the new equilibrium have:</u>

	FORWARD RATE	REVERSE RATE
A.	increased	increased
B.	not changed	increased
C.	decreased	increased
D.	not changed	not changed

85 E2 Consider the following equilibrium:

$$H_{2(g)} + I_{2(g)} \rightleftharpoons 2HI_{(g)}$$

The volume of the equilibrium system is increased and a new equilibrium is established. <u>Compared to the rates in the original equilibrium</u>, which of the following describes the rates of the forward and reverse reactions in the new equilibrium?

FORWARD RATE	REVERSE RATE
decreased	decreased
increased	increased
decreased	increased
remained constant	remained constant

86 E3 Consider the following equilibrium:

$$4\text{HCl}_{(g)} + \text{O}_{2(g)} \rightleftharpoons 2\text{H}_2\text{O}_{(g)} + 2\text{Cl}_{2(g)} + \text{energy}$$

The temperature of the equilibrium system is <u>increased</u> and a <u>new equilibrium is established</u>. The rates of the forward and reverse reactions for the new equilibrium <u>compared to the original equilibrium</u> have

	FORWARD RATE	REVERSE RATE
A.	increased	increased
B.	decreased	not changed
C.	decreased	increased
D.	not changed	increased

87 E3 Consider the following equilibrium reaction:

$$PCl_{5(g)} + energy \rightleftharpoons PCl_{3(g)} + Cl_{2(g)}$$

The temperature of this system is decreased. What is the immediate effect on the reaction rates?

- A. Both forward and reverse rates increase.
- B. Both forward and reverse rates decrease.
- C. Forward rate decreases while reverse rate increases.
- D. Forward rate increases while reverse rate decreases.
- An equilibrium system shifts left when the temperature is increased. The forward reaction is A. exothermic and ΔH is positive.
   C. endothermic and ΔH is positive.
   D. endothermic and ΔH is negative.

### 89 E3 An equilibrium system shifts <u>left</u> when the

- A. rate of the forward reaction is equal to the rate of the reverse reaction.
- B. rate of the forward reaction is less than the rate of the reverse reaction.
- C. rate of the forward reaction is greater than the rate of the reverse reaction.
- D. rate of the forward reaction and the rate of the reverse reaction are constant.

### 90 E4 Addition of a <u>catalyst</u> to an equilibrium system

- A. increases the value of  $K_{eq}$ . B. increases the yield of products.
- C. has no effect on the rates of reaction. D. increases the rate of formation of both reactants and products.

31	E4	Consider the following equilibrium:				
		$2SO_{2(g)} + O_{2(g)} \rightleftharpoons 2SO_{3(g)} \qquad \Delta H = -19$	8 kJ			
		There will be no shift in this equilibrium when				
		A. more $O_2$ is added. B. a catalyst is a	dded.			
92	F4	C. the volume is increased. D. the temperate	ure is ir	ncreased.		
52	<u> </u>	$2SO \pm O \rightarrow 2SO \qquad AH = -107 \text{ kJ}$				
		$230_{2(g)} + 0_{2(g)} \leftarrow 230_{3(g)} \qquad \Delta \Pi = -197 \text{ KJ}$				
		Which of the following will <u>not</u> shift the equilibriu	m to t	he right?		
93	E4	A. adding more $O_2$ B. adding a catalyst C. Incr A catalyst is added to a system already at	easing	the pressure L	. lowering the temperat	ure
50		equilibrium. How are the forward and reverse		FORWARD RATE	REVERSE RATE	
		reaction rates affected by the <u>addition of the</u> catalyst?	A.	increases	increases	
			В.	increases	remains constant	
			C.	remains constant	decreases	
			D.	remains constant	remains constant	
01	F5	Ethene C.H. can be produced in the following indus	trial ev	etem:		
54	20		that by			
		$C_2H_{6(g)}$ + energy $\rightleftarrows$ $C_2H_{4(g)}$ + $H_{2(g)}$				
		The conditions that are necessary to maximize theA. low temperature and low pressure.B. low	<u>e equi</u> tempe	librium yield of ( rature and high pr	<u>2₂H₄</u> are essure.	
95	E5	C. high temperature and low pressure. D. high	tempe	erature and high p	ressure.	
		C. high temperature and low pressure.D. highConsider the following equilibrium:	tempe	erature and high p	ressure.	
		C. high temperature and low pressure.D. highConsider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + energy$	tempe	K <sub>eq</sub>	ressure. Equilibrium Position	
		C. high temperature and low pressure. D. high Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + energy$	A.	K <sub>eq</sub>	RESSURE. EQUILIBRIUM POSITION favours products	
		C. high temperature and low pressure. D. high Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + energy$ Certain conditions provide less than 10% yield of NH <sub>2</sub> at equilibrium. Which of the following	A. B.	K <sub>eq</sub> large small	EQUILIBRIUM POSITION favours products favours products	
		C. high temperature and low pressure. D. high Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + energy$ Certain conditions provide less than 10% yield of NH <sub>3</sub> at equilibrium. Which of the following describes this equilibrium?	A. B. C.	Keq     Iarge       small     Iarge	EQUILIBRIUM POSITION favours products favours reactants	
		C. high temperature and low pressure. D. high Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + energy$ Certain conditions provide less than 10% yield of NH <sub>3</sub> at equilibrium. Which of the following describes this equilibrium?	A. B. C. D.	K <sub>eq</sub> Iarge       small     Iarge	EQUILIBRIUM POSITION favours products favours products favours reactants favours reactants	
96	E5	C. high temperature and low pressure. D. high Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + energy$ Certain conditions provide less than 10% yield of NH <sub>3</sub> at equilibrium. Which of the following describes this equilibrium? Consider the following equilibrium system:	A. B. C. D.	K <sub>eq</sub> Iarge       small     Iarge	EQUILIBRIUM POSITION favours products favours products favours reactants favours reactants	
96	E5	C. high temperature and low pressure. D. high Consider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + energy$ Certain conditions provide less than 10% yield of NH <sub>3</sub> at equilibrium. Which of the following describes this equilibrium? Consider the following equilibrium system: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + 92 \text{ kJ}$	A. B. C. D.	K <sub>eq</sub> I       large     small       large     small	EQUILIBRIUM POSITION favours products favours products favours reactants favours reactants	
96	E5	C. high temperature and low pressure.D. highConsider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + energy$ Certain conditions provide less than 10% yield of NH3 at equilibrium. Which of the following describes this equilibrium?Consider the following equilibrium?Consider the following equilibrium system: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + 92 \text{ kJ}$ Which of the following sets of conditions will fav. A. low pressure and low temperatureB. low D. high pressure and low temperatureD. high	A. B. C. D. D.	Keq       Keq         large       small         large       small         small       small         small       formation of the sure and high tem ssure and	EQUILIBRIUM POSITION favours products favours products favours reactants favours reactants favours reactants	
96	E5	C. high temperature and low pressure.D. highConsider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + energy$ Certain conditions provide less than 10% yield of NH3 at equilibrium. Which of the following describes this equilibrium?Consider the following equilibrium?Consider the following equilibrium system: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + 92 \text{ kJ}$ Which of the following sets of conditions will fave A. low pressure and low temperature Consider the following equilibrium:A. low pressure and low temperature C. high pressure and low temperature D. high Consider the following equilibrium:	A. B. C. D. D.	Keq       Iarge         large       small         large       small         small       Iarge         Iarge       Iarge         Iarg	EQUILIBRIUM POSITION favours products favours reactants favours reactants favours reactants product?	
96 97	E5	C. high temperature and low pressure.D. highConsider the following equilibrium: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + energy$ Certain conditions provide less than 10% yield of NH3 at equilibrium. Which of the following describes this equilibrium?Consider the following equilibrium system: $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + 92 \text{ kJ}$ Which of the following sets of conditions will fave A. low pressure and low temperature C. high pressure and low temperature D. high Consider the following equilibrium: $Cl_2O_{7(g)} + 8H_{2(g)} \rightleftharpoons 2HCl_{(g)} + 7H_{2(g)}$	A. B. C. D. D. D. D. D. D. D. D. D. D	Keq         Iarge         small         large         small         large         small         large         small         large         small         large         small         large         small         small         g)	EQUILIBRIUM POSITION favours products favours reactants favours reactants favours reactants product? perature nperature	

**98** E5 Consider the following equilibrium:

99

# $2NO_{2(g)} \rightleftharpoons N_2O_{4(g)} + energy$

### The number of moles of NO<sub>2</sub> at equilibrium could be increased by

A. adding  $N_2O_4$  B. adding a catalyst.

C. decreasing the temperature D. decreasing the volume by increasing the pressure.

E5 Ammonia, NH<sub>3</sub>, is produced by the following reaction:

$$N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + energy$$

Which of the following would result in the highest concentration of ammonia at equilibrium?

A. increasing the temperature and increasing the pressure

B. decreasing the temperature and increasing the pressure

C. increasing the temperature and decreasing the pressure

D. decreasing the temperature and decreasing the pressure

**100** E5 Methanol,  $CH_3OH$ , can be produced by the following:

$$CO_{(g)} + 2H_{2(g)} \rightleftharpoons CH_3OH_{(g)} + energy$$

### The conditions that are necessary to maximize the equilibrium yield of CH<sub>3</sub>OH are

A. low temperature and low pressure.	B. high temperature and low pressure.
C. low temperature and high pressure.	D. high temperature and high pressure.

### THE EQUILIBRIUM CONSTANT

### 101 F1 Which of the following reactions most favours products?

	REACTION	K <sub>eq</sub>	A. I	B. II	C. III
Ι	$2\operatorname{SO}_{2(g)} + \operatorname{O}_{2(g)} \rightleftharpoons 2\operatorname{SO}_{3(g)}$	$2.6 \times 10^2$			
Π	$2 \operatorname{NO}_{(g)} + \operatorname{O}_{2(g)} \rightleftharpoons 2 \operatorname{NO}_{2(g)}$	$6.4 \times 10^5$			
Ш	$2\operatorname{CO}_{(g)} + \operatorname{O}_{2(g)} \rightleftharpoons 2\operatorname{CO}_{2(g)}$	$2.5 \times 10^{15}$			
IV	$2 \operatorname{H}_{2(g)} + \operatorname{O}_{2(g)} \rightleftharpoons 2 \operatorname{H}_{2} \operatorname{O}_{(g)}$	$1.7 \times 10^{27}$			

102	F1	An indication t	hat an equilibrium sy	stem <u>favours the products</u>	is a
		A. large K <sub>eq</sub> .	B. positive ΔH.	C. one step mechanism.	D. low activation energy.
103	F1	Which of the fo	llowing reactions mo	ost <u>favors the reactants</u> ?	
		A. $CH_{4(g)} \in$	$2 H_{2(g)} + C_{(s)}$	$K_{eq} = 1.2 \times 10^{-9}$	
		B. $SbCl_{5(g)}$	$\rightleftharpoons$ SbCl <sub>3(g)</sub> + Cl <sub>2(g)</sub>	$K_{eq} = 2.5 \times 10^{-2}$	
		C. $N_2O_{4(g)}$	$\rightleftharpoons$ 2NO <sub>2(g)</sub>	$K_{eq} = 4.5 \times 10^{-1}$	
		D. $C_{(s)} + CO_{s}$	$_{2(g)} \rightleftharpoons 2\mathrm{CO}_{(g)}$	$K_{eq} = 1.4 \times 10^1$	

AUG 2000

D. IV

104 F1 Which equation has the largest value of  $K_{eq}$ ? A.  $N_{2(g)} + O_{2(g)} \rightleftharpoons 2NO_{(g)} \qquad \Delta H = 21 \text{ kJ}$ B.  $C_2H_{6(g)} \rightleftharpoons 2C_{(g)} + 3H_{2(g)}$  $\Delta H = 83 \text{ kJ}$ C.  $H_{2(g)} + \frac{1}{2}O_{2(g)} \rightleftharpoons H_2O_{(g)}$  $\Delta H = -240 \text{ kJ}$ D.  $\operatorname{Ca}_{(s)} + 2\operatorname{H}_2\operatorname{O}_{(\ell)} \rightleftharpoons \operatorname{Ca}(\operatorname{OH})_{2(aq)} + \operatorname{H}_{2(g)} \qquad \Delta \operatorname{H} = -240 \text{ kJ}$ **105** F1 Consider the following equilibrium:  $K_{eq} = 2.1 \times 10^{30}$  $2 \operatorname{NO}_{(\sigma)} \rightleftharpoons \operatorname{N}_{2(\sigma)} + \operatorname{O}_{2(\sigma)}$ The value of the equilibrium constant indicates that the A.  $[NO]^2 < [N_2][O_2]$ B.  $[NO]^2 > [N_2][O_2]$ C.  $[NO] = [N_2][O_2]$ D.  $[NO] > [N_2][O_2]$ 106 F1 Which of the following equilibrium systems most <u>favours the products</u>?  $K_{ea} = 6.4 \times 10^{-39}$ A.  $\operatorname{Cl}_{2(q)} \rightleftharpoons \operatorname{2Cl}_{(q)}$ B.  $\operatorname{Cl}_{2(g)} + 2\operatorname{NO}_{(g)} \rightleftharpoons 2\operatorname{NOCl}_{(g)}$   $K_{eq} = 3.7 \times 10^8$ C.  $\operatorname{Cl}_{2(g)} + 2\operatorname{NO}_{2(g)} \rightleftharpoons 2\operatorname{NO}_2\operatorname{Cl}_{(g)} \quad K_{eq} = 1.8$  $K_{eq} = 2.0 \times 10^{-7}$ D.  $2 \text{HCl}_{(g)} \rightleftharpoons \text{H}_{2(g)} + \text{Cl}_{2(g)}$ **107** F1 Consider the following equilibrium:  $\operatorname{COCl}_{2(g)} \rightleftharpoons \operatorname{CO}_{(g)} + \operatorname{Cl}_{2(g)}$ At equilibrium in a 1.0 L container, there are 3.0 mol COCl<sub>2</sub>, 0.49 mol CO and 0.49 mol Cl<sub>2</sub>. At constant temperature the volume of the above system is decreased to 0.50 L. When equilibrium is reestablished the

A. concentrations of all three gases have increased.

B. concentrations of all three gases have decreased.

C.  $[COCl_2]$  has increased and [CO] and  $[Cl_2]$  have decreased.

D. [COCl<sub>2</sub>] has decreased and [CO] and [Cl<sub>2</sub>] have increased.

**108** F1 Consider the following equilibrium:

2

$$\operatorname{NOCl}_{(g)} \rightleftharpoons 2\operatorname{NO}_{(g)} + \operatorname{Cl}_{2(g)}$$

In a 1.0 L container at equilibrium there are 1.0 mol NOCI, 0.70 mol NO and 0.40 mol Cl<sub>2</sub>.

	[NOC1]	[NO]	[Cl <sub>2</sub> ]
A.	new = old	new = old	new = old
B.	new > old	new > old	new > old
C.	new < old	new < old	new > old
D.	new < old	new > old	new > old

At constant temperature and volume, 0.10 mol NOCI is added. The concentrations in the <u>"new"</u> equilibrium in <u>comparison</u> to the

concentrations in the <u>"old"</u> equilibrium are

**109** F1 Consider the following equilibrium:

$$2 H_2 O_{(g)} \rightleftharpoons 2 H_{2(g)} + O_{2(g)}$$

When 0.1010 mol  $H_2O$  is placed in a 1.000 L container, equilibrium is established. The equilibrium concentration of  $O_2$  is 0.0010 mol/L. The equilibrium concentrations of  $H_2O$  and  $H_2$ are

	[H <sub>2</sub> O]	[H <sub>2</sub> ]
Α.	0.0990	0.0020
В.	0.1000	0.0010
C.	0.1005	0.0005
D.	0.1010	0.0020

AUG 2000

110 F1 Consider the following equilibrium:  

$$COCl_{2(g)} \rightleftharpoons CO_{(g)} + Cl_{2(g)} \qquad K_{eq} = 8.1 \times 10^{-4}$$
For the above system.....  
For the above system.....  

$$COCl_{2(g)} \rightleftharpoons CO_{(g)} + Cl_{2(g)} \qquad K_{eq} = 8.1 \times 10^{-4}$$

$$COCl_{2} = [CO][Cl_{2}]$$

$$C. \quad [COCl_{2}] = [CO][Cl_{2}]$$

$$C. \quad [COCl_{2}] = \frac{1}{[CO][Cl_{2}]}$$

**111** F1 Consider the following equilibrium:

$$2O_{2(g)} + N_{2(g)} \rightleftharpoons N_2O_{4(g)}$$

When 2.0 mol of  $O_2$  and 3.0 mol of  $N_2$  were placed in a 10.0 L container at 25° C, the value of  $K_{eq}$  = 0.90. If the same number of moles of reactant were placed in a 5.0 L container at 25° C, the <u>equilibrium constant</u> would be

**A.** 
$$0.011$$
 **B.**  $0.45$  **C.**  $0.90$  **D.**  $1.80$   
**112** F1 Consider the following equilibrium system at 900° C:

$$H_2O_{(g)} + CO_{(g)} \rightleftharpoons H_{2(g)} + CO_{2(g)}$$

Initially 5.0 moles of  $H_2O$  and 4.0 moles of CO were reacted. At equilibrium, it is found that2.0 moles of  $H_2$  are present. How many moles of  $H_2O$  remain in the mixture?A. 1.0 molesB. 2.0 molesC. 3.0 molesD. 4.0 moles

$$\mathrm{CO}_{2(g)} + \mathrm{H}_{2(g)} \rightleftharpoons \mathrm{CO}_{(g)} + \mathrm{H}_{2}\mathrm{O}_{(g)}$$

1.00 mole of  $CO_2$  and 2.00 moles of  $H_{2(g)}$  are placed into a 2.00 litre container. At equilibrium, the [CO] = 0.31 mol/L. **Based on this data, the equilibrium** [CO<sub>2</sub>] is A. 0.19 M B. 0.31 M C. 0.38 M D. 0.69 M

**114** F1 Consider the following equilibrium:

$$H_{2(g)} + I_{2(g)} \rightleftharpoons 2HI_{(g)} K_{eq} = 50.0$$

What is the <u>value</u>  $K_{eq}$  for the reaction rewritten as:

$$2HI_{(g)} \rightleftharpoons H_{2(g)} + I_{2(g)} K_{eq} = ?$$
  
A. -50.0 B. 0.0200 C. 25.0 D. 50.0

115	F2	Consider the following equilibrium:	
		$2\mathrm{Fe}_{(s)} + 3\mathrm{H}_2\mathrm{O}_{(g)} \rightleftharpoons \mathrm{Fe}_2\mathrm{O}_{3(s)} + 3\mathrm{H}_{2(g)}$	
		The equilibrium constant expression is A. $K_{eq} = \frac{[Fe_2O_3][H_2]^3}{[Fe]^2[H_2O]^3}$ B.	$K_{eq} = \frac{[Fe_2O_3][3H_2]}{[2Fe][3H_2O]}$
		C. $K_{eq} = \frac{[H_2]^3}{[H_2O]^3}$ D.	$\mathbf{K}_{eq} = \left[\mathbf{H}_2\right]^3$
116	F2	Consider the following equilibrium:	
		$2H_2S_{(g)} \rightleftharpoons 2H_{2(g)} + S_{2(g)}$	Δ (0.10)(0.40)
		At equilibrium, $[H_2S]$ =0. 50 mol/L, $[H_2]$ =0.10 mol/L and $[S_2]$ =0. 40 mol/L.	(0.50)
		The value of $K_{eq}$ is calculated using the ratio	B. $\frac{(0.10)^2(0.40)}{(0.50)^2}$
			C. $\frac{(0.50)}{(0.10)(0.50)}$
			D. $\frac{(0.50)^2}{(0.10)^2(0.40)}$
117	F2	For which of the following equilibria does $K_{eq} = [O_2]$ ?	
		A. $O_{2(l)} \rightleftharpoons O_{2(g)}$	
		B. $2O_{3(q)} \rightleftharpoons 3O_{2(q)}$	
		C. $2H_2O_{(l)} \rightleftharpoons 2H_{2(g)} + O_{2(g)}$	
		D. $2 \text{Hg}_{(s)} + \text{O}_{2(g)} \rightleftharpoons 2 \text{HgO}_{(s)}$	
118	F2	<ul> <li>Which of the following statements is <u>correct</u>?</li> <li>A. K<sub>eq</sub> is the ratio of [products] to [reactants].</li> <li>B. K<sub>eq</sub> determines how fast a reaction is completed.</li> <li>C. A large K<sub>eq</sub> value indicates that reactants are favoured.</li> <li>D. A small K<sub>eq</sub> value indicates that products are favoured.</li> </ul>	
119	F2	Consider the following equilibrium system:	
		$3O_{2(g)} \rightleftharpoons 2O_{3(g)}$ $K_{eq} = 1$ Which equation compares the concentration of oxygen and ozone?	A. $[O_2] = [O_3]^{\frac{2}{3}}$
		the concentration of oxygen and ozone i	B. $[O_2] = [O_3]$
			C. $[O_2] = [O_3]^{\frac{3}{2}}$
			D. $[O_2]^{\frac{2}{3}} = [O_3]$
-			

120	F2	F2 An equal number of moles of $I_{2(g)}$ and $Br_{2(g)}$ are placed into a closed container and allowed to establish the following equilibrium:					
		$I_{2(g)} + Br_{2(g)} \rightleftharpoons 2IBr_{(g)} \qquad K_{eq} = 2$	280 A	. $[I_2] = [IBr]$			
	Which one of the following <u>relates</u> [IBr] to [	2] at B	. [I <sub>2</sub> ]<[IBr]				
		equilibrium?	С	. $[I_2] = 2[IBr]$			
			D	. $[I_2] = 280[IBr]$			
121	F2	Consider the following reaction:					
		$2 \text{Hg}_{(g)} + \text{O}_{2(g)} \rightleftharpoons 2 \text{HgO}_{(s)}$	А	$K_{eq} = \frac{1}{\left[\mathrm{Hg}\right]^2 \left[\mathrm{O}_2\right]}$			
		The equilibrium constant expression for the	e reaction is B	$K_{eq} = [Hg]^2 [O_2]$			
			С	$K_{eq} = \frac{\left[\text{HgO}\right]^2}{\left[\text{Hg}\right]^2 \left[\text{O}_2\right]}$			
			D	$K_{eq} = \frac{[2 \text{HgO}]}{[2 \text{Hg}][\text{O}_2]}$			
122	F2	Consider the following reaction:					
		$2H_{2(g)} + O_{2(g)} \rightleftharpoons 2H_2O_{(l)}$		A. $K_{eq} = [H_2]^2 [O_2]$			
		What is the equilibrium constant expression	<u>n</u> for the reaction?	B. $K_{eq} = \frac{[H_2]^2[O_2]}{[H_2O]^2}$			
				C. $K_{eq} = \frac{[H_2O]^2}{[H_2]^2[O_2]}$			
				D. $K_{eq} = \frac{1}{\left[H_2\right]^2 \left[O_2\right]}$			
123	F2	Consider the following equilibrium:	[\$0	,]			
		$2 \operatorname{SO}_{2(g)} + \operatorname{O}_{2(g)} \rightleftharpoons 2 \operatorname{SO}_{3(g)}$	A. $K_{eq} = \frac{100}{[SO_2][}$	0 <sub>2</sub> ]			
		The equilibrium expression is:	B. $K_{eq} = \frac{[SO_2]^2}{[SO_2]^2}$	$\begin{bmatrix} O_2 \end{bmatrix}$			
			C. $K_{eq} = \frac{[SO_2][}{[SO_2]}$	$O_2]$			
			D. $K_{eq} = \frac{[SO_2]^2}{[SO_3]}$	$\left[ \underbrace{O_2}_{l^2} \right]$			



**127** F2 Consider the following equilibrium:

$$2N_2O_{(g)} + 3O_{2(g)} \rightleftharpoons 4NO_{2(g)}$$

The equilibrium constant expression is...

A. 
$$K_{eq} = \frac{[2N_2O][3O_2]}{[4NO_2]}$$
  
B.  $K_{eq} = \frac{[N_2O]^2[O_2]^3}{[NO_2]^4}$   
C.  $K_{eq} = \frac{[4NO_2]}{[2N_2O][3O_2]}$   
D.  $K_{eq} = \frac{[NO_2]^4}{[N_2O]^2[O_2]^3}$ 

128 F2 Given the following equilibrium system:

 $\operatorname{Br}_{2(g)} \rightleftharpoons \operatorname{Br}_{2(1)}$ A.  $K_{eq} = \frac{\left[Br_{2(1)}\right]}{\left[Br_{2(q)}\right]}$ The equilibrium constant expression for the above system is..... B.  $K_{eq} = \left[ Br_{2(g)} \right]$ C.  $K_{eq} = \frac{1}{\left[Br_{2(q)}\right]}$ D.  $K_{eq} = \left[ Br_{2(g)} \right] \left[ Br_{2(g)} \right]$ F2 Consider the following equilibrium: 129  $4\mathrm{KO}_{2(s)} + 2\mathrm{H}_2\mathrm{O}_{(g)} \rightleftharpoons 4\mathrm{KOH}_{(s)} + 3\mathrm{O}_{2(g)}$ The equilibrium constant A.  $K_{eq} = \frac{[KOH]^4 [O_2]^3}{[KO_2]^4 [H_2O]^2}$  B.  $K_{eq} = \frac{[O_2]^3}{[H_2O]^2}$ expression is..... C.  $K_{eq} = \frac{[KO_2]^4 [H_2O]^2}{[KOH]^4 [O_2]^3}$  D.  $K_{eq} = \frac{[H_2O]^2}{[O_2]^3}$ F2 Consider the following equilibrium: 130 A.  $\frac{[2CO]^2[O_2]}{[2CO_2]^2}$  $2CO_{(g)} + O_{2(g)} \rightleftharpoons 2CO_{2(g)}$ The ratio used to calculate the equilibrium constant is B.  $\frac{[2CO_2]^2}{[2CO]^2[O_2]}$ C.  $\frac{[CO]^2[O_2]}{[CO_2]^2}$ D.  $\frac{\left[CO_{2}\right]^{2}}{\left[CO\right]^{2}\left[O_{2}\right]}$ 131 F2 Consider the following equilibrium: A.  $K_{eq} = [H^+][I^-]$  $I_{2(s)} + H_2O_{(\ell)} \rightleftharpoons H^+_{(aa)} + I^-_{(aa)} + HOI_{(aa)}$ B.  $K_{eq} = [H^+][I^-][HOI]$ The equilibrium constant expression for the above system is..... C.  $K_{eq} = \frac{\left[H^+\right]\left[I^-\right]\left[HOI\right]}{\left[I_2\right]\left[H_2O\right]}$ D.  $K_{eq} = \frac{\left[H^+\right]\left[I^-\right]\left[HOI\right]}{\left[H_2O\right]}$  132 F2 The equilibrium constant expression for the reaction below is....

		$2 Hg_{(l)} + O_{2(g)} \rightleftharpoons 2 HgO_{(s)}$			A.	$\mathbf{K}_{eq} = \frac{1}{\left[\mathbf{O}_{2}\right]}$
					B.	$\mathbf{K}_{eq} = \begin{bmatrix} \mathbf{O}_2 \end{bmatrix}$
					C.	$K_{eq} = \frac{[2HgO]}{[O_2][2Hg]}$
					D.	$\mathbf{K}_{eq} = \frac{\left[\mathrm{HgO}\right]^2}{\left[\mathrm{Hg}\right]^2 \left[\mathrm{O}_2\right]}$
133	F2	Consider the following equilibrium system:				
		$\operatorname{SnO}_{2(s)} + 2\operatorname{CO}_{(g)} \rightleftharpoons \operatorname{Sn}_{(s)} + 2\operatorname{CO}_{2(g)}$			А.	$K_{eq} = \frac{[CO_2]^2}{[CO]^2}$
		The <u>equilibrium constant expression</u> for the al	bove	e system is	В.	$K_{eq} = \frac{[2CO_2]^2}{[2CO]^2}$
					C.	$K_{eq} = \frac{[CO_2]^2 [Sn]}{[CO]^2 [SnO_2]}$
					D.	$K_{eq} = \frac{[2CO_2]^2 [Sn]}{[2CO]^2 [SnO_2]}$
134	F2	Consider the following equilibrium constant expres	ssior	1:		
		$K_{eq} = [CO_2]$	A.	$\operatorname{CO}_{2(g)} \rightleftharpoons \mathbf{C}$	$CO_{2(s)}$	)
		Which one of the following <u>equilibrium</u> <u>systems</u> does the above <u>expression</u> represent?	В.	$PbO_{(s)} + CO_2$	2(g) <del>(</del>	$\neq$ PbCO <sub>3(s)</sub>
			C.	$CaCO_{3(s)} \rightleftharpoons$	CaO	$C_{(s)} + CO_{2(g)}$
			D.	$H_2CO_{3(aq)}$	≓ H <sub>2</sub>	$_{2}O_{(l)} + CO_{2(aq)}$
135	F2	What is the $\underline{K}_{eq}$ expression for		AUG 2000		[, ] <sup>2</sup>
		$\mathrm{Sb}^{3+}_{(aq)} + \mathrm{Cl}^{-}_{(aq)} + \mathrm{H}_2\mathrm{O}_{(\ell)} \rightleftharpoons \mathrm{SbOCl}_{(s)} + 2\mathrm{I}_{(s)}$	$\mathrm{H}^{+}_{(aq)}$	1)	Α.	$\mathbf{K}_{eq} = \frac{\left[\mathbf{H}^{+}\right]}{\left[\mathbf{Sb}^{3+}\right]\left[\mathbf{Cl}^{-}\right]}$
					В.	$\mathbf{K}_{eq} = \frac{\left[\mathbf{H}^{+}\right]^{2} \left[\mathbf{SbOCl}\right]}{\left[\mathbf{Sb}^{3+}\right] \left[\mathbf{Cl}^{-}\right]}$
					C.	$K_{eq} = \frac{\left[H^{+}\right]^{2}}{\left[Sb^{3+}\right]\left[Cl^{-}\right]\left[H_{2}O\right]}$

D.  $K_{eq} = \frac{\left[H^+\right]^2 \left[SbOCI\right]}{\left[Sb^{3+}\right] \left[CI^-\right] \left[H_2O\right]}$ 

**136** F2 The equilibrium expression for a reaction is.....

The reaction could be:

A. 
$$6H^{+}_{(aq)} + BiS_{(s)} \rightleftharpoons 2Bi^{3+}_{(aq)} + 3H_2S_{(g)}$$
  
B.  $6H^{+}_{(aq)} + Bi_2S_{3(s)} \rightleftharpoons 2Bi^{3+}_{(aq)} + 3H_2S_{(g)}$   
C.  $2Bi^{3+}_{(aq)} + 3H_2S_{(aq)} \rightleftharpoons Bi_2S_{3(s)} + 6H^{+}_{(aq)}$   
D.  $2Bi^{3+}_{(aq)} + 3H_2S_{(aq)} \rightleftharpoons Bi_2S_{3(aq)} + 6H^{+}_{(aq)}$ 

137 F2

<sup>2</sup> What is the  $K_{eq}$  expression for the following reaction?

$$\operatorname{SnO}_{2(s)} + 2\operatorname{CO}_{(g)} \rightleftharpoons \operatorname{Sn}_{(s)} + 2\operatorname{CO}_{2(g)}$$

A. 
$$K_{eq} = \frac{[CO_2]}{[CO]}$$
  
B.  $K_{eq} = \frac{[CO_2]^2}{[CO]^2}$   
C.  $K_{eq} = \frac{[Sn][CO_2]^2}{[CO]^2}$   
D.  $K_{eq} = \frac{[Sn][CO_2]^2}{[SnO_2][CO]^2}$ 

138 F2 What is the  $K_{eq}$  expression for the following equilibrium?

$$3Fe_{(s)} + 4H_{2}O_{(g)} \rightleftharpoons Fe_{3}O_{4(s)} + 4H_{2(g)} \qquad A. \quad K_{eq} = [H_{2}]^{4}$$

$$B. \quad K_{eq} = \frac{[H_{2}]}{[H_{2}O]}$$

$$C. \quad K_{eq} = \frac{[H_{2}]^{4}}{[H_{2}O]^{4}}$$

$$D. \quad K_{eq} = \frac{[Fe_{3}O_{4}][H_{2}]^{4}}{[Fe]^{3}[H_{2}O]^{4}}$$

 $\mathbf{K}_{eq} = \frac{\left[\mathbf{H}^{+}\right]^{6}}{\left[\mathbf{B}\mathbf{i}^{3+}\right]^{2}\left[\mathbf{H}_{2}\mathbf{S}\right]^{3}}$ 

**139** F3 Consider the following equilibrium:

$$2NO_{(g)} + 2H_{2(g)} \rightleftharpoons N_{2(g)} + 2H_2O_{(g)} \qquad K_{eq} = 1.3 \times 10^2$$

A 1.0 L container is initially filled with 1.0 mol of each of the species in the reaction. <u>The equilibrium shifts</u> to the:

A. left because Trial  $K_{eq} > K_{eq}$ B. left because Trial  $K_{eq} < K_{eq}$ C. right because Trial  $K_{eq} > K_{eq}$ D. right because Trial  $K_{eq} < K_{eq}$ 

### 140 F3 Identify the equilibrium system that least favors the formation of products.

A.	$2 \text{HgO}_{(s)} \rightleftharpoons 2 \text{Hg}_{(l)} + \text{O}_{2(g)}$	$K_{eq} = 1.2 \times 10^{-22}$
В.	$\mathrm{CH}_{3}\mathrm{COOH}_{(aq)} + \mathrm{H}_{2}\mathrm{O}_{(l)} \rightleftharpoons \mathrm{H}_{3}\mathrm{O}_{(aq)}^{+} + \mathrm{CH}_{3}\mathrm{COO}_{(aq)}^{-}$	$K_{eq} = 1.8 \times 10^{-5}$
C.	$2 \operatorname{NO}_{(g)} + \operatorname{O}_{2(g)} \rightleftharpoons 2 \operatorname{NO}_{2(g)}$	$K_{eq} = 6.5 \times 10^5$
D.	$H_{2(g)} + Cl_{2(g)} \rightleftharpoons 2HCl_{(g)}$	$K_{eq} = 1.8 \times 10^{33}$



149 F4 Consider the following equilibrium:  $CaCO_{3(s)} + 556 \text{ kJ} \rightleftharpoons CaO_{(s)} + CO_{2(g)}$ The value of the equilibrium constant will increase when A. CO<sub>2</sub> is added. B. CO<sub>2</sub>is removed. C. the temperature is increased. D. the temperature is decreased. F4 Consider the following equilibrium: 150  $2 \operatorname{NO}_{(g)} + \operatorname{Cl}_{2(g)} \rightleftharpoons 2 \operatorname{NOCl}_{(g)}$ At constant temperature and volume, Cl<sub>2</sub> is added to the above equilibrium system. As equilibrium reestablishes, the B. K<sub>eq</sub> will decrease. C. [NO] will increase. A. K<sub>eq</sub> will increase. D. [NOCI] will increase. 151 F4 In an exothermic equilibrium reaction involving only gases, the value of K<sub>eq</sub> can be <u>decreased</u> by A. adding some reactant gas. B. removing some reactant gas. C. increasing the temperature. D. decreasing the temperature. F4 Consider the following equilibrium system: 152  $CO_{(q)} + 2H_{2(q)} \rightleftharpoons CH_3OH_{(q)}$  $\Delta H = -18 \text{ kJ}$ In order to increase the value of Keq for this reaction, you could A. increase [CO] B. increase the volume C. decrease [CH<sub>3</sub>OH] D. decrease the temperature F4 The temperature of an exothermic reaction at equilibrium is increased by 10° C. The value of Keq 153 A. doubles B. increases C. decreases D. remains constant. F4 Consider the following equilibrium: 154  $\operatorname{Co}(\operatorname{H}_2\operatorname{O})^{2+}_{6\ (aa)}$  +  $4\operatorname{Cl}^-_{(aq)}$   $\rightleftharpoons$   $\operatorname{Co}\operatorname{Cl}^{2-}_{4\ (aq)}$  +  $6\operatorname{H}_2\operatorname{O}_{(\ell)}$ (pink) (blue) When the temperature is increased, the solution turns a dark blue. Based on this observation, the reaction is A. exothermic and the  $K_{eq}$  has increased. B. exothermic and the  $K_{eq}$  has decreased. C. endothermic and the  $K_{eq}$  has increased. D. endothermic and the  $K_{eq}$  has decreased. F4 155 Consider the following reaction:  $\Delta H = -74.8 \, kJ$  $C_{(s)} + 2H_{2(g)} \rightleftharpoons CH_{4(g)}$ Which of the following will cause an <u>increase</u> in the value of  $K_{eq}$ ? A. increasing [H<sub>2</sub>] B. decreasing the volume C. finely powdering the  $C_{(s)}$ D. decreasing the temperature **156** F4 Consider the following equilibrium:  $PCl_{5(g)} \rightleftharpoons PCl_{3(g)} + Cl_{2(g)}$  $\Delta H = +92.5 \text{ kJ}$ When the temperature decreases, the equilibrium A. shifts left and K<sub>eq</sub> value increases. B. shifts left and K<sub>eq</sub> value decreases. C. shifts right and K<sub>eq</sub> value increases. D. shifts right and Keq value decreases F4 Consider the following equilibrium: 157  $\Delta H = +181 \text{ kJ}$  $N_{2(g)} + O_{2(g)} \rightleftharpoons 2NO_{(g)}$ When the temperature is decreased, the equilibrium A. shifts left and the  $K_{eq}$  value increases. B. shifts left and the K<sub>eq</sub> value decreases. C. shifts right and the  $K_{eq}$  value increases. D. shifts right and the  $K_{eq}$  value decreases



**165** F5 Consider the following equilibrium:

$$H_{2(g)} + I_{2(g)} \rightleftharpoons 2HI_{(g)}$$

At equilibrium the  $[H_2] = 0.020 \text{ mol/L}$ ,  $[I_2] = 0.020 \text{ mol/L}$  and [HI] = 0.160 mol/L.

The value of the equilibrium constant is...

A. 
$$2.5 \times 10^{-3}$$
 B.  $1.6 \times 10^{-2}$  C.  $6.4 \times 10^{1}$  D.  $4.0 \times 10^{2}$ 

**166** F5 Consider the following equilibrium:

$$C_{(s)} + H_2O_{(g)} \rightleftharpoons CO_{(g)} + H_{2(g)}$$

The contents of a 1.00 L container at equilibrium were analyzed and found to contain 0.20 mol C, 0.20 mol H<sub>2</sub>O, 0.60 mol CO and 0.60 mol H<sub>2</sub>. The <u>equilibrium constant</u> is..

	A. 0.11	B. 0.56	C. 1.8	D. 9.0	
F5	Consider the follo	wing equilibrium:			

**167** F5 Consider the following equilibrium:

$$H_{2(g)} + S_{(s)} \rightleftharpoons H_2 S_{(g)}$$

In a 1.0 L container at equilibrium there are 0.050 mol  $H_2$  , 0.050 mol S and 1.0 mol  $H_2S$ . The value of  $K_{e\sigma}$  is

A.  $2.5 \times 10^{-3}$  B.  $5.0 \times 10^{-2}$  C.  $2.0 \times 10^{1}$  D.  $4.0 \times 10^{2}$ 

**168** F5 Consider the following system and concentrations at equilibrium:

What is the value of  $K_{eq}$  for the above system?

$$2NO_{(g)} + Br_{2(g)} \rightleftharpoons 2NOBr_{(g)}$$

Substance	Equilibrium Concentration
NO	$1.2 \times 10^{-2} \text{ mol/L}$
$Br_2$	$3.4 \times 10^{-2}$ mol/L
NOBr	$5.8 \times 10^{-1} \text{ mol/L}$

**169** F5 Consider the following equilibrium system:

A. 1.5 X10<sup>-5</sup>

C. 1.4 X10<sup>3</sup>

$$\operatorname{PCl}_{5(g)} \rightleftharpoons \operatorname{PCl}_{3(g)} + \operatorname{Cl}_{2(g)}$$

B. 8.2 X<sup>1</sup>0<sup>2</sup>

D. 6.9 X10<sup>4</sup>

At equilibrium, [PCl<sub>5</sub>] is 0.400M, [PCl<sub>3</sub>] is 1.50M and [Cl<sub>2</sub>] is 0.600M. The K<sub>eq</sub> for the reaction is

$$2$$
SO<sub>2(g)</sub> + O<sub>2(g)</sub>  $\rightleftharpoons$  2SO<sub>3(g)</sub>

At equilibrium,  $[SO_2]$  is 4.00 x10<sup>-3</sup> mol/L,  $[O_2]$  is 4.00x10<sup>-3</sup> mol/L and  $[SO_3]$  is 2.33x10<sup>-3</sup> mol/L. From this data, **the K<sub>eq</sub> value for the above system is** 

171 F5 Consider the following equilibrium system:

 $\operatorname{CO}_{(g)} + \operatorname{Cl}_{2(g)} \rightleftharpoons \operatorname{COCl}_{2(g)}$ 

At equilibrium, a 2.0 litre sample was found to contain 1.00 mol CO, 0.500 mol  $Cl_2$  and 0.100 mol  $COCl_2$ . The  $K_{eq}$  value for the above system is A. 0.40 B. 0.20 C. 2.5 D. 5.0  $H_{2(g)} + I_{2(g)} \rightleftharpoons 2HI_{(g)}$ 

At equilibrium  $[H_2] = 0.00220 \text{ mol/L}, [I_2] = 0.00220 \text{ mol/L} \text{ and } [HI] = 0.0156 \text{ mol/L}.$ The value of Keq is

A.  $3.10 \times 10^{-4}$ 

B.  $1.99 \times 10^{-2}$ 

- C.  $5.03 \times 10^{1}$
- D.  $3.22 \times 10^3$

**173** F6 Consider the following equilibrium:

$$\operatorname{PCl}_{3(g)} + \operatorname{Cl}_{2(g)} \rightleftharpoons \operatorname{PCl}_{5(g)}$$

When 0.40 mol of PCI<sub>3</sub> and 0.40 mol of CI<sub>2</sub> are placed in a 1.00 L container and allowed to reach equilibrium, 0.244 mol of PCI<sub>5</sub> are present. From this information, the value of Keg is

B. 0.30 C. 3.3 A. 0.10 D. 10 174 F6 Consider the following:

 $2C_{(s)} + O_{2(g)} \rightleftharpoons 2CO_{(g)}$ 

A 1.00 L flask is initially filled with 2.00 mol C and 0.500 mol  $O_2$ . At equilibrium, the  $[O_2]$ is 0.250 mol/L. The K<sub>eq</sub> value is

		A. 0.444	B. 1.00	C. 2.00	D. 2.25	
175	F7	Consider the following equilibrium:				

175 Consider the following equilibrium: F/

$$2 \operatorname{HBr}_{(g)} \rightleftharpoons \operatorname{H}_{2(g)} + \operatorname{Br}_{2(g)}$$

Initially, 0.100 mol HBr is placed into a 2.0 L container. At equilibrium, there are 0.040 mol HBr present. The equilibrium concentration of H<sub>2</sub> is

A. 0.0050 mol/L C. 0.015 mol/L D. 0.030 mol/L B. 0.010 mol/L **176** F7 Consider the following equilibrium:

 $N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)}$ 

A 1.00 L container is initially filled with 0.200 mol  $N_2O_4$ . At equilibrium, 0.160 mol  $NO_2$  are present. What is the equilibrium concentration of N2O4?

		A. 0. 040 mol/L	B. 0. 080 mol/L	C. 0.120 mol/L	D. 0.160 mol/L	
177	F7	Consider the followi	ng equilibrium:			
		$2 \operatorname{NOBr}_{(g)} \rightleftharpoons$	$2\operatorname{NO}_{(g)} + \operatorname{Br}_{2(g)}$	$K_{eq} = 6.4 \times 10^7$	-2	
		At equilibrium, a 1	.00 L flask contains (	0.030 mol NOBr and	0.030 mol NO.	
		How many mol Br <sub>2</sub>	are present?			
		A. 1.9 x10 <sup>-3</sup> mol	B. 6.4 x10 <sup>-2</sup> mol	C. 3.0 x10 <sup>-2</sup> mol	D. 4.7 x10 <sup>-1</sup> mol	

178	F7 *	Consider the following equilibrium: $CO_{(g)} + 2H_{2(g)} \rightleftharpoons CH_3OH_{(g)} \qquad K_{eq}$	= 12.0			
		At equilibrium, a 1.00 L flask contains 0.020 mol CO and What is the <u>concentration of <math>CH_3OH</math> at equilibrium?</u>	0.35 mol $H_2$ .			
179	F7	A. $2.0 \times 10^{-4}$ mol/L B. $5.8 \times 10^{-4}$ mol/L C. $2.9 \times 10^{-2}$ r Consider the following equilibrium:	mol/L D. 8.4 x 10 <sup>-2</sup> mol/L			
		$N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)} \qquad K_{eq} = 1.0 \times 10^{-2}$				
		At equilibrium, the $[NO_2] = 2.0 \times 10^{-2} \text{ mol/L}$ and the $[N_2O_2] = 2.0 \times 10^{-2} \text{ mol/L}$	D <sub>4</sub> ] is			
		A. 4.0x10 <sup>-6</sup> mol/L B. 4.0x10 <sup>-2</sup> mol/L C. 2.0 mol/L	D. 25 mol/L	_		
180	F7 *	Consider the following equilibrium:				
		$\operatorname{CH}_{4(g)} + \operatorname{H}_2\operatorname{O}_{(g)} \rightleftharpoons \operatorname{CO}_{(g)} + 3\operatorname{H}_{2(g)} \qquad \operatorname{K}_{eq} = 5.7$				
		At equilibrium, the $[CH_4] = 0.40 \text{ mol/L}, [CO] = 0.30 \text{ mol/L} \text{ and } [H_2]$ The $[H_2O]$ is	= 0.80  mol/L.			
		A. 0.067 mol/L				
		B. 0.11 mol/L C. 2.2 mol/L				
		D. 5.3 mol/L				
181	F7 *	Consider the following equilibrium: $N \cap \rightarrow 2NO \qquad K = 4.61 \times 10^{-3}$				
		$N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)} \qquad K_{eq} = 4.61 \times 10^{-3}$				
		A 1.00 L container at equilibrium was analyzed and for At equilibrium, the concentration of $N_2O_4$ is	ound to contain 0.0200 mol N	10 <sub>2</sub> .		
182	F7	A 1.00 L container at equilibrium was analyzed and for At <u>equilibrium</u> , the concentration of $N_2O_4$ is A. 0. 0868 mol/L B. 0. 230 mol/L C. 4.34 mol Consider the following equilibrium:	ound to contain 0.0200 mol N /L D. 11.5 mol/L	10 <sub>2</sub> .		
182	F7 *	A 1.00 L container at equilibrium was analyzed and for At <u>equilibrium</u> , the concentration of N <sub>2</sub> O <sub>4</sub> is A. 0. 0868 mol/L B. 0. 230 mol/L C. 4.34 mol Consider the following equilibrium: $N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)}$ $K_{eq} = 0.133$	ound to contain 0.0200 mol № /L D. 11.5 mol/L	IO <sub>2</sub> .		
182	F7 *	A 1.00 L container at equilibrium was analyzed and for At <u>equilibrium</u> , the concentration of N <sub>2</sub> O <sub>4</sub> is A. 0. 0868 mol/L B. 0. 230 mol/L C. 4.34 mol Consider the following equilibrium: $N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)}$ $K_{eq} = 0.133$ At equilibrium, the [N <sub>2</sub> O <sub>4</sub> ] is equal to	ound to contain 0.0200 mol N /L D. 11.5 mol/L	IO <sub>2</sub> .		
182	F7 *	A 1.00 L container at equilibrium was analyzed and for At <u>equilibrium</u> , the concentration of N <sub>2</sub> O <sub>4</sub> is A. 0. 0868 mol/L B. 0. 230 mol/L C. 4.34 mol Consider the following equilibrium: $N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)}$ $K_{eq} = 0.133$ <u>At equilibrium</u> , the [N <sub>2</sub> O <sub>4</sub> ] is equal to	A. $\frac{0.133}{[NO_2]}$	$B. = \frac{[NO_2]}{0.133}$		
182	F7 *	A 1.00 L container at equilibrium was analyzed and for At <u>equilibrium</u> , the concentration of N <sub>2</sub> O <sub>4</sub> is A. 0. 0868 mol/L B. 0. 230 mol/L C. 4.34 mol Consider the following equilibrium: $N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)}$ $K_{eq} = 0.133$ At equilibrium, the [N <sub>2</sub> O <sub>4</sub> ] is equal to	A. $\frac{0.133}{[NO_2]}$	$B. = \frac{[NO_2]}{0.133}$		
182	F7 *	A 1.00 L container at equilibrium was analyzed and for At <u>equilibrium</u> , the concentration of N <sub>2</sub> O <sub>4</sub> is A. 0. 0868 mol/L B. 0. 230 mol/L C. 4.34 mol Consider the following equilibrium: $N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)}$ $K_{eq} = 0.133$ At equilibrium, the [N <sub>2</sub> O <sub>4</sub> ] is equal to	A. $\frac{0.133}{[NO_2]}$ C. $\frac{0.133}{[NO_2]^2}$	<b>IO<sub>2</sub></b> . B. $\frac{[NO_2]}{0.133}$ D. $\frac{[NO_2]^2}{0.133}$		
182	F7 * F7	A 1.00 L container at equilibrium was analyzed and for At <u>equilibrium</u> , the concentration of N <sub>2</sub> O <sub>4</sub> is A. 0. 0868 mol/L B. 0. 230 mol/L C. 4.34 mol Consider the following equilibrium: $N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)}$ $K_{eq} = 0.133$ At equilibrium, the [N <sub>2</sub> O <sub>4</sub> ] is equal to	Dund to contain 0.0200 mol N         /L       D. 11.5 mol/L         A. $\frac{0.133}{[NO_2]}$ C. $\frac{0.133}{[NO_2]^2}$	<b>IO</b> <sub>2</sub> . B. $\frac{[NO_2]}{0.133}$ D. $\frac{[NO_2]^2}{0.133}$		
182	F7 * F7	A 1.00 L container at equilibrium was analyzed and for At <u>equilibrium</u> , the concentration of N <sub>2</sub> O <sub>4</sub> is A. 0. 0868 mol/L B. 0. 230 mol/L C. 4.34 mol Consider the following equilibrium: $N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)}$ $K_{eq} = 0.133$ At equilibrium, the [N <sub>2</sub> O <sub>4</sub> ] is equal to Consider the following equilibrium: $2NO_{(g)} + Cl_{2(g)} \rightleftharpoons 2NOCl_{(g)}$ $K_{eq} = 12$	A. $0.133 \\ [NO_2]$ C. $0.133 \\ [NO_2]^2$	$HO_2$ . B. $\frac{[NO_2]}{0.133}$ D. $\frac{[NO_2]^2}{0.133}$		
182	F7 *	A 1.00 L container at equilibrium was analyzed and for At equilibrium, the concentration of N2O4 is A. 0. 0868 mol/LB. 0. 230 mol/LC. 4.34 molConsider the following equilibrium:N2O4(g) $\rightleftharpoons$ 2NO2(g)Keq = 0.133At equilibrium, the [N2O4] is equal toConsider the following equilibrium:2NO(g) + Cl2(g) $\rightleftharpoons$ 2NOCl(g)Keq = 12At equilibrium, [NOCI]=1.60 mol/L and [NO]=0.80 mol/LAt equilibrium, [NOCI]=1.60 mol/LC. 0. 33 mol/L	A. $0.133 \\ [NO_2]$ C. $0.133 \\ [NO_2]^2$ Z. $0.133 \\ [NO_2]^2$ D. $0.133 \\ [NO_2]^2$	$HO_2$ . B. $\frac{[NO_2]}{0.133}$ D. $\frac{[NO_2]^2}{0.133}$		
182 183 184	F7 * F7 *	A 1.00 L container at equilibrium was analyzed and for At equilibrium, the concentration of N <sub>2</sub> O <sub>4</sub> is A. 0. 0868 mol/L B. 0. 230 mol/L C. 4.34 mol Consider the following equilibrium: $N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)}$ $K_{eq} = 0.133$ At equilibrium, the [N <sub>2</sub> O <sub>4</sub> ] is equal to Consider the following equilibrium: $2NO_{(g)} + Cl_{2(g)} \rightleftharpoons 2NOCl_{(g)}$ $K_{eq} = 12$ At equilibrium, [NOCI]=1.60 mol/L and [NO]=0.80 mol/L A. 0.17 mol/L B. 0. 27 mol/L C. 0. 33 mol/L Consider the following equilibrium system:	A. $0.133 \\ [NO_2]$ C. $0.133 \\ [NO_2]^2$ Z.       The [Cl <sub>2</sub> ] is D. 3. 0 mol/L	$HO_2$ . B. $\frac{[NO_2]}{0.133}$ D. $\frac{[NO_2]^2}{0.133}$		
182 183 184	F7 * F7 *	A 1.00 L container at equilibrium was analyzed and for At <u>equilibrium</u> , the concentration of N <sub>2</sub> O <sub>4</sub> is A. 0. 0868 mol/L B. 0. 230 mol/L C. 4.34 mol Consider the following equilibrium: $N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)}$ $K_{eq} = 0.133$ At equilibrium, the [N <sub>2</sub> O <sub>4</sub> ] is equal to Consider the following equilibrium: $2NO_{(g)} + Cl_{2(g)} \rightleftharpoons 2NOCl_{(g)}$ $K_{eq} = 12$ At equilibrium, [NOCI]=1.60 mol/L and [NO]=0.80 mol/L A. 0.17 mol/L B. 0. 27 mol/L C. 0. 33 mol/L Consider the following equilibrium system: $2NO_{(g)} + O_{2(g)} \rightleftharpoons 2NO_{2(g)}$ $K_{eq} = 6$	A. $0.133 \\ [NO_2]$ C. $0.133 \\ [NO_2]^2$ <b>/L. The [Cl<sub>2</sub>] is</b> D. 3. 0 mol/L	$HO_2$ . B. $\frac{[NO_2]}{0.133}$ D. $\frac{[NO_2]^2}{0.133}$		
182 183 184	F7 * F7 *	A 1.00 L container at equilibrium was analyzed and for At equilibrium, the concentration of N <sub>2</sub> O <sub>4</sub> is A. 0. 0868 mol/L B. 0. 230 mol/L C. 4.34 mol Consider the following equilibrium: $N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)}$ $K_{eq} = 0.133$ At equilibrium, the [N <sub>2</sub> O <sub>4</sub> ] is equal to Consider the following equilibrium: $2NO_{(g)} + Cl_{2(g)} \rightleftharpoons 2NOCl_{(g)}$ $K_{eq} = 12$ At equilibrium, [NOCI]=1.60 mol/L and [NO]=0.80 mol/L A. 0.17 mol/L B. 0. 27 mol/L C. 0. 33 mol/L Consider the following equilibrium system: $2NO_{(g)} + O_{2(g)} \rightleftharpoons 2NO_{2(g)}$ $K_{eq} = 6$ At equilibrium, the [NO] = 0.600 M and the [O <sub>2</sub> ] = 0.300 M [NO <sub>2</sub> ] is: A. 7.0 M B. 3.4 M C. 2.6 M D. 0	Jund to contain 0.0200 mol N         /L       D. 11.5 mol/L         A. $\frac{0.133}{[NO_2]}$ C. $\frac{0.133}{[NO_2]^2}$ /L. The [Cl <sub>2</sub> ] is D. 3. 0 mol/L         55         1. Using this data, the equilit .60 M	$HO_2 .$ B. $\frac{[NO_2]}{0.133}$ D. $\frac{[NO_2]^2}{0.133}$		

Consider the following equilibrium: 185

AUG 2000

$$2NO_{2(g)} \rightleftharpoons N_2O_{4(g)} \qquad K_{eq} = 1.15$$

The equilibrium concentration of NO<sub>2</sub> is 0.50 mol/L. Calculate the equilibrium concentration of N<sub>2</sub>O<sub>4(g)</sub>

A. 0.22 mol/LB. 0.29 mol/LConsider the following equilibrium: C. 0.43mol/L D. 0.58mol/L 186

$$2O_{3(g)} \rightleftharpoons 3O_{2(g)} \qquad K_{eq} = 36$$

What is the concentration of  $O_3$  when the equilibrium concentration of  $O_2$  is  $6.0 \times 10^{-2}$  mol/L?

- A.  $2.4 \times 10^{-3} \text{ mol/L}$
- B.  $4.0 \times 10^{-2} \text{ mol/L}$
- C.  $6.0 \times 10^{-2} \text{ mol/L}$
- D.  $9.0 \times 10^{-2} \text{ mol/L}$

Consider the following equilibrium: F8 187

$$PCl_{5(g)} \rightleftharpoons PCl_{3(g)} + Cl_{2(g)} \qquad K_{eq} = 2.30$$

A 1.0 L container is filled with 0.05 mol  $PCI_5$ , 1.0 mol  $PCI_3$ , and 1.0 mol  $CI_2$ .

### The system proceeds to the

A. left because Trial  $K_{eq} > K_{eq}$ B. left because Trial  $K_{eq} < K_{eq}$ C. right because Trial  $K_{eq} > K_{eq}$ D. right because Trial  $K_{eq} < K_{eq}$ F8Consider the following equilibrium:

188

$$H_2O_{(g)} + Cl_2O_{(g)} \rightleftharpoons 2HOCl_{(g)} \qquad K_{eq} = 9.0 \times 10^{-2}$$

A 1.0 L flask contains a mixture of 1.8  $\times 10^{-1}$  mol H<sub>2</sub>O, 4.0  $\times 10^{-4}$  mol Cl<sub>2</sub>O, and 8.0 $\times 10^{-2}$  mol HOCI. To establish equilibrium, the system will proceed to the

A. left because Trial  $K_{eq} > K_{eq}$ B. left because Trial  $K_{eq} < K_{eq}$ C. right because Trial  $K_{eq} > K_{eq}$ D. right because Trial  $K_{eq} < K_{eq}$ 

189 F8 Consider the following equilibrium:

 $2O_{3(g)} \rightleftharpoons 3O_{2(g)} \qquad K_{eq} = 55$ 

If 0.060 mol of O<sub>3</sub> and 0.70 mol of O<sub>2</sub> are introduced into a 1.0 L vessel, the

- A.  $K_{trial} > K_{eq}$  and the  $[O_2]$  increases.
- B.  $K_{trial} < K_{eq}$  and the  $[O_2]$  increases.

C. 
$$K_{trial} > K_{eq}$$
 and the  $[O_2]$  decreases.

D.  $K_{trial} < K_{eq}$  and the  $[O_2]$  decreases.

**190** F8 Consider the following equilibrium:

$$N_{2(g)} + O_{2(g)} \rightleftharpoons 2NO_{(g)} \qquad K_{eq} = 0.010$$

<u>Initially</u>, a 1.0 L container is filled with 0.40 mol of  $N_2$ , 0.10 mol of  $O_2$  and 0.080 mol of NO. As the system <u>approaches</u> equilibrium the

A. [NO], [N<sub>2</sub>] and [O<sub>2</sub>] remain unchanged.

B. [NO] increases and both [N<sub>2</sub>] and [O<sub>2</sub>] decrease.

C. [NO] decreases and both  $[N_2]$  and  $[O_2]$  increase.

D. [NO] decreases and both [N<sub>2</sub>] and [O<sub>2</sub>] remain unchanged.

**191** F8 Consider the following:

$$2 \operatorname{NO}_{2(g)} \rightleftharpoons \operatorname{N}_2 \operatorname{O}_{4(g)} \qquad \operatorname{K}_{eq} = 1.20$$

A 1.0 L flask is filled with 1.4 mol  $NO_2$  and 2.0 mol  $N_2O_4$  . To reach equilibrium, the reaction proceeds to the

A. left as Trial  $K_{eq} > K_{eq}$ B. left as Trial  $K_{eq} < K_{eq}$ C. right as Trial  $K_{eq} > K_{eq}$ D. right as Trial  $K_{eq} < K_{eq}$ 

**192** F8 Consider the following equilibrium:

$$PCl_{5(g)} \rightleftharpoons PCl_{3(g)} + Cl_{2(g)} \qquad K_{eq} = 33.3$$

<u>Predict what will occur</u> when 2.0 mol of  $PCI_5$ , 3.0 mol of  $PCI_3$  and 4.0 mol of  $CI_2$  are placed in a 1.0 L container and allowed to establish equilibrium.

A. [PCl₅] will increase	B. [PCl <sub>3</sub> ] and [Cl <sub>2</sub> ] will both increase
C. [PCl <sub>5</sub> ] and [Cl <sub>2</sub> ] will both increase	D. [PCl <sub>5</sub> ] and [ PCl <sub>3</sub> ] will both decrease

**193** F8 Consider the following equilibrium system:

 $2SO_{2(g)} + O_{2(g)} \rightleftharpoons 2SO_{3(g)} \qquad K_{eq} = 4.0$ 

In an experiment, 0.40 mol  $SO_{2(g)}$ , 0.20 mol  $O_{2(g)}$  and 0.40 mol  $SO_{3(g)}$  are placed

into a 1.0 litre container. Which of the following statements relates the changes in

 $[SO_2]$  and  $[O_2]$  as equilibrium becomes established?

A. The $[SO_2]$ and $[O_2]$ increase.	B. The $[SO_2]$ and $[O_2]$ decrease.
C. The [SO <sub>2</sub> ] and $[O_2]$ do not change	D. The $[SO_2]$ increases and the $[O_2]$ decreases.

194 F8

AUG 2000

Consider the following equilibrium:

 $2\text{NOCl}_{(g)} \rightleftharpoons 2\text{NO}_{(g)} + \text{Cl}_{2(g)}$ 

A flask is filled with NOCl, NO and  $Cl_2$ . Initially there was a total of 5.0 moles of gases present. When equilibrium is reached, there is a total of 6.0 moles of gases present. Which of the following explains this observation?

- A. The reaction proceeded left because the Trial  $K_{eq} > K_{eq}$
- B. The reaction proceeded left because the Trial  $K_{eq} < K_{eq}$
- C. The reaction proceeded right because the Trial  $K_{eq} > K_{eq}$
- D. The reaction proceeded right because the Trial  $K_{eq} < K_{eq}$

**195** F8 Consider the following equilibrium:

$$2\text{NOCl}_{(g)} \rightleftharpoons 2\text{NO}_{(g)} + \text{Cl}_{2(g)}$$

A flask of fixed volume is initially filled with NOCl<sub>(g)</sub>, NO<sub>(g)</sub>, and Cl<sub>2(g).</sub> When equilibrium is reached, the pressure has increased. To reach equilibrium, the reaction proceeded to the A. left because Trial K<sub>eq</sub> was less than K<sub>eq</sub> B. right because Trial K<sub>eq</sub> was less than K<sub>eq</sub> C. left because Trial K<sub>eq</sub> was greater than K<sub>eq</sub> D. right because Trial K<sub>eq</sub> was greater than K<sub>eq</sub>

### **ANSWERS TO MULTIPLE CHOICE QUESTIONS:**

INTROD	UCTION				
1.	В	26. D	PRINCIPLE	73. A	98. A
2.	Α	27. C	49. A	74. A	99. B
3.	С	28. D	50. C	75. D	100. C
4.	С	29. C	51. C	76. C	
5.	С	30. C	52. C	77. C	THE EQUILIBRIUM
6.	D	31. C	53. C	78. A	CONSTANT
7.	В	32. D	54. B	79. C	101. D
8.	С	33. B	55. C	80. C	102. A
9.	С	34. C	56. A	81. B	103. A
10.	D	35. A	57. B	82. D	104. D
11.	Α	36. C	58. B	83. C	105. A
12.	В	37. D	59. B	84. A	106. B
13.	D	38. B	60. B	85. A	107. A
14.	С	39. A	61. B	86. A	108. B
15.	Α	40. B	62. A	87. B	109. A
16.	С	41. A	63. A	88. B	110. C
17.	D	42. D	64. D	89. B	111. C
18.	В	43. A	65. D	90. D	112. C
19.	В	44. D	66. D	91. B	113. A
20.	D	45. C	67. A	92. B	114. B
21.	С	46. A	68. A	93. A	115. C
22.	В	47. C	69. B	94. C	116. B
23.	D	48. C	70. C	95. D	117. A
24.	D		71. B	96. C	118. A
25.	D	LE	72. A	97. B	119. A
		CHATELIER'S			120. B

G DYNAMIC EQUILIBRIUM STUDY GUIDE 2000

Page 34 of 52

121. A	136. C	151. C	166. C	181. A
122. D	137. B	152. D	167. C	182. D
123. B	138. C	153. C	168. D	183. C
124. B	139. D	154. C	169. D	184. C
125. B	140. A	155. D	170. C	185. B
126. C	141. B	156. B	171. A	186. A
127. D	142. C	157. B	172. C	187. A
128. C	143. B	158. B	173. D	188. A
129. B	144. B	159. B	174. B	189. C
130. D	145. D	160. B	175. C	190. C
131. B	146. C	161. A	176. C	191. D
132. A	147. D	162. A	177. B	192. B
133. A	148. B	163. D	178. C	193. A
134. C	149. C	164. B	179. B	194. D
135. A	150. D	165. C	180. A	195. B

### DYNAMIC EQUILIBRIUM STUDY GUIDE 2000

### written

### INTRODUCTION TO EQUILIBRIUM

### D3 1 Consider the following equilibrium:

$$2 \operatorname{NOCl}_{(g)} \rightleftharpoons 2 \operatorname{NO}_{(g)} + \operatorname{Cl}_{2(g)}$$

A chemist places 2.00 mol NOCI in a 1.0 L container. Describe the changes in [NOCI] and  $[Cl_2]$  as the system approaches equilibrium. (**1mark**)

D4	2	Identify four characteristics of a chemical equilibrium. (2 marks)
D4	3	What is "equal" in a chemical reaction that has reached a state of equilibrium? 2 marks
D5	4	a) Why are chemical equilibria referred to as dynamic? (1 mark)
		b) How is a chemical system at equilibrium recognized? (1 mark)
D7	5	Consider the following equilibrium:
		$4\text{HCl}_{(g)} + \text{O}_{2(g)} \rightleftharpoons 2\text{H}_2\text{O}_{(g)} + 2\text{Cl}_{2(g)} + \text{energy}$

a) How does the entropy change in the forward direction? Explain your reasoning. (1 mark)b) How does the enthalpy change in the forward direction? Explain your reasoning. (1 mark)

D9 6 **Describe how enthalpy and entropy change**, in the forward direction, as an exothermic reaction reaches equilibrium. **Explain your reasoning**. (2 marks)

### LE CHATELIER'S PRINCIPLE

- E1 7 State Le Chatelier's Principle. (2 marks)
- E1 8 State Le Chatelier's Principle. (2 marks)

E2 9 Consider the following equilibrium:

$$2 \operatorname{NO}_{(g)} + \operatorname{Cl}_{2(g)} \rightleftharpoons 2 \operatorname{NOCl}_{(g)} \qquad \Delta \mathrm{H} = -77 \text{ kJ}$$

What happens to the amount of  $Cl_2$  when the following changes are imposed?

Explain, using Le Chatelier's principle.

a) Removing NO<sub>(g)</sub> . (1 mark)

b) Decreasing the temperature. (1 mark)

E2 10 Consider the following equilibrium:

$$CO_{(g)} + 2H_{2(g)} \rightleftharpoons CH_3OH_{(g)} \qquad \Delta H = -18 \text{ kJ}$$

Explain, using Le Chatelier's principle, how the following changes will affect the number of moles of  $CH_3OH$  present at equilibrium.

a) Adding a catalyst. (1 mark)

b) Decreasing the volume of the system. (1 mark)

E2 11 Consider the following equilibrium:

$$PCl_{3(g)} + Cl_{2(g)} \rightleftharpoons PCl_{5(g)} \qquad \Delta H = -88 \text{ kJ}$$

What happens to the  $[PCl_3]$  when additional  $Cl_2$  is added at constant temperature and volume? Explain. **(2marks)** 

E2 12 Consider the following equilibrium:

$$2\operatorname{CrO_4}^{2-}_{(aq)} + \operatorname{H_2O}_{(\ell)} \rightleftharpoons \operatorname{Cr_2O_7}^{2-}_{(aq)} + 2\operatorname{OH}_{(aq)}^{-}$$
vellow orange

When HCl is added drop-by-drop to the yellow solution above, the solution turns orange. Explain why this colour change occurs. **(2marks)** 

E2 13 Consider the following equilibrium:

$$N_2H_{4(g)} + 2O_{2(g)} \rightleftharpoons 2NO_{(g)} + 2H_2O_{(g)}$$

More oxygen is added to the above equilibrium. After the system re-establishes equilibrium, identify the substance(s), if any, that have a net (2 marks)

a) increase in concentration.

b) decrease in concentration.

E2 14 Consider the following equilibrium system:

 $\operatorname{Fe}_{(aq)}^{3+} + \operatorname{SCN}_{(aq)}^{-} \rightleftharpoons \operatorname{FeSCN}_{(aq)}^{2+}$ yellow colourless red

In an experiment, a student places the above equilibrium system into a cold water bath and notes that the intensity of the red colour increases. The student then concludes that the equilibrium is exothermic.

a) Do you agree or disagree? 0.5 mark

b) Explain: **1.5 marks** E2 15 Consider the following

$$\operatorname{Fe}^{3+}_{(aq)}$$
 +  $\operatorname{SCN}^{-}_{(aq)}$   $\rightleftharpoons$   $\operatorname{FeSCN}^{2+}_{(aq)}$   
yellow colourless red

When a few drops of 6.0 M NaOH is added to 25.0 mL of the above system, a precipitate of Fe(OH)<sub>3</sub> forms and the solution turns pale yellow.

- a) Explain this colour change in terms of Le Chatelier's Principle. (2marks)
- b) Describe the effect on the rate of the reverse reaction as the colour change
- occurs. (1mark)

E2 16 Consider the observations for the following equilibrium:

$$N_2O_{4(g)} \rightleftharpoons 2NO_{2(g)}$$

(colourless) (brown)

Trial	Temperature °C	Colour
I.	10	light brown
п.	50	dark brown

a) Sketch the potential energy curve on the graph below for this equilibrium. (1mark)



b) Explain the colour change using Le Chytelier's Principle. (1mark)

c) Other than changing temperature, what could be done to cause a shift to the left? (1mark)

E2 17 Methanol, CH<sub>3</sub>OH, is produced industrially by the following reaction:

$$CO_{(g)} + 2H_{2(g)} \rightleftharpoons CH_3OH_{(g)} + heat$$

a) State two different methods of shifting the equilibrium to the right. (1 mark)

b) In terms of rates, explain why these methods cause the equilibrium to shift to the right. (1mark)

E3 18 Consider the following equilibrium:

$$2H_2O_{(g)} \rightleftharpoons 2H_{2(g)} + O_{2(g)}$$

Identify two ways to increase the rate of the forward reaction. (2marks)

E3 19 Consider the following equilibrium:

HInd +  $H_2O \rightleftharpoons H_3O^+$  +  $Ind^-$ 

(yellow)

(blue)

The system is yellow and turns blue on the addition of NaOH. In terms of the forward and reverse reaction rates, explain why this shift occurs. **(2marks)** 

### THE EQUILIBRIUM CONSTANT

F1 20 Consider the following equilibrium system:

$$N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + energy$$

A 1.00 L container is filled with 5.0 mol  $NH_3$  and the system proceeds to equilibrium as

indicated by the graph.



- a) Draw and label the graph for  $N_2$  and  $H_2$ . (2 marks)
- b) Calculate the  $K_{eq}$  for  $N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)}$ . (2 marks)

F1 21 Consider the following equilibrium:

$$H_{2(g)} + I_{2(g)} \rightleftharpoons 2HI_{(g)} \qquad K_{eq} = 64$$

Equal moles of  $H_2$  and  $I_2$  are placed in a 1.00 L container. At equilibrium, the [HI]=0.160 mol/L. Calculate the initial [ $H_2$ ]. (3 marks)

F1 22 Consider the following equilibrium:



 $H_{2(g)} + I_{2(g)} \rightleftharpoons 2HI_{(g)}$ 

A 2.0L container is filled with 0.070 mol of  $H_2$  and 0.060mol of  $I_2$ . Equilibrium is reached after 15.0 minutes at which time there is 0.060 mol of HI present. Sketch and label the graphs for the changes in

concentrations of  $H_2$ ,  $I_2$ , and HI for the time period of 0 to 30.0 minutes. **(3marks)** 

F2 23 Consider the following equilibrium:

$$2 \operatorname{NO}_{(g)} + \operatorname{O}_{2(g)} \rightleftharpoons 2 \operatorname{NO}_{2(g)} \qquad \operatorname{K}_{eq} = 6.45 \times 10^5$$

a) Write the  $K_{eq}$  expression. (1mark)

b) Explain why the [NO<sub>2</sub>] is greater than the [NO] at equilibrium when the [O<sub>2</sub>] is 1.0 mol/L. (1mark)

24 Consider the following equilibrium: F4

$$CS_{2(g)} + 3Cl_{2(g)} \rightleftharpoons CCl_{4(g)} + S_2Cl_{2(g)} \qquad \Delta H = -238 \text{ kJ}$$

- a) Sketch a potential energy diagram for the reaction above and label ΔH. (2 marks)
- b) Some CS<sub>2</sub> is added and equilibrium is then reestablished. State the direction of the equilibrium shift and the resulting change in [Cl<sub>2</sub>]. (1 mark)
- c) The temperature is decreased and equilibrium is then reestablished. What will the effect be on the value of K<sub>eq</sub> ? (1 mark)
- F5 25 Consider the graph below representing the following equilibrium:

$$CH_3CH_2CH_2CH_{3(g)} \rightleftharpoons CH_3CH(CH_3)_{2(g)}$$

n-butane isobutane

Data for the graph was obtained from various equilibrium mixtures.

Calculate the value of  $K_{eq}$  for the equilibrium. (2 marks)



26 F5 Consider the following diagram for a chemical system containing three substances



represented by A, B and C:

- a) What feature of the graph indicates that the system reaches equilibrium? (1 mark)
- b) Write a balanced equation for the equilibrium reaction. (2 marks)
- c) Calculate K<sub>eq</sub> at equilibrium. (2 marks)

reactants PE Progress of the reaction

F5 27 Consider the following graph for the reaction: energy  $\square + N_2O_4 \square_{(g)} \square \Rightarrow 2NO_2 \square_{(g)}$ 



- a) What is the stress imposed at time  $t_1$  ? (1mark)
- b) What is the stress imposed at time  $t_3$  ? (1mark)
- c) Calculate  $K_{eq}$  for the equilibrium between  $t_2$  and  $t_3$  . (2marks)

F5 28 Consider the following equilibrium:

$$2NO_{(g)} + O_{2(g)} \rightleftharpoons 2NO_{2(g)}$$

At 227° C in a 2.00 L container there are 0.044 mol NO, 0.100 mol  $O_2$ 

and 7.88 mol NO<sub>2</sub> at equilibrium. Calculate the equilibrium constant. (3marks)

F6 29 At high temperature, 0.500 mol HBr was placed in a 1.00 L container where it decomposed to give the equilibrium:

$$2 \operatorname{HBr}_{(g)} \rightleftharpoons \operatorname{H}_{2(g)} + \operatorname{Br}_{2(g)}$$

At equilibrium, the [Br<sub>2</sub>] is 0.0855 mol/L. What is the value of the equilibrium constant? (**3 marks**) F6 30 Consider the following equilibrium:

$$2\mathrm{CO}_{2(g)} ~\rightleftarrows~ 2\mathrm{CO}_{(g)} + \mathrm{O}_{2(g)}$$

Initially, a 1.0 L container is filled with 0.050 mol of  $CO_2$ . At equilibrium, the  $[CO_2]$  is 0.030 mol/L. Calculate the value of  $K_{eq}$ . (3marks)

F6 31 Consider the following equilibrium:

$$2 \operatorname{CH}_{4(g)} \rightleftharpoons \operatorname{C}_2 \operatorname{H}_{2(g)} + 3 \operatorname{H}_{2(g)}$$

A 0.180 mol sample of  $CH_4$  is added to an empty 1.00 L container. At equilibrium,

the [C<sub>2</sub>H<sub>2</sub>] is 0.0800 mol/L. Calculate the equilibrium constant. (4 marks)

F6 32 In an experiment, 0.200 mol of CO (g) and 0.400 mol of O<sub>2(g)</sub> are placed in a 1.00 L container and the following equilibrium is achieved:

$$2\mathrm{CO}_{(g)} + \mathrm{O}_{2(g)} \rightleftharpoons 2\mathrm{CO}_{2(g)}$$

At equilibrium, the  $[CO_2]$  is found to be 0.160 mol/L. Calculate the value of  $K_{eq}$ . **(3marks)** 33 Given the following equilibrium:

$$H_{2(g)} + I_{2(g)} \rightleftharpoons 2HI_{(g)}$$

Initially, 0.200 mol  $H_2$  and 0.200 mol  $I_2$  were placed into a 1.0 L container. At equilibrium, the [ $I_2$ ] is 0.040 mol/L. Calculate the  $K_{ea}$ . (3 marks)

F6 34 Consider the following equilibrium system:

$$\mathrm{PCl}_{3(g)} + \mathrm{Cl}_{2(g)} \rightleftharpoons \mathrm{PCl}_{5(g)}$$

At 250°C, 0.40 mol of PCl<sub>3</sub> and 0.60 mol of Cl<sub>2</sub> are placed into a 1.0 litre container. At equilibrium, the [PCl<sub>5</sub>] = 0.11 mol/L. Calculate the value of  $K_{eq}$ . (3 marks)

F6

### F6 35 Consider the following equilibrium: AUG 2000

 $3I_{2(g)} + 3F_{2(g)} \rightleftharpoons 2IF_{2(g)} + I_4F_{2(g)}$ 

Initially,  $2.00 \times 10^{-1}$  mol of  $I_2$  and  $3.00 \times 10^{-1}$  mol of  $F_2$  are put into a 10.00 L flask. At equilibrium,  $[I_4F_2]$  is  $2.00 \times 10^{-3}$  M. Calculate the  $K_{eq}$ . (4 marks)

F6 36 Consider the following equilibrium:

 $\operatorname{Fe}_{(aa)}^{3+} + \operatorname{SCN}_{(aa)}^{-} \rightleftharpoons \operatorname{FeSCN}_{(aa)}^{2+}$ 

Initially, 50.0 mL of  $0.10 \text{ M Fe}^{3+}$  is added to 30.0 mL of  $0.20 \text{ M SCN}^{-}$ . At equilibrium, the concentration of FeSCN<sup>2+</sup> is found to be 0.050 M. Calculate the K<sub>ea</sub> for the reaction.

F7 37 Consider the following equilibrium:

$$H_{2(g)} + I_{2(g)} \rightleftharpoons 2HI_{(g)} \qquad K_{eq} = 1.2 \times 10^{-2}$$

A 2.0 L flask is filled with 0.10 mol HI. Calculate the concentration of H<sub>2</sub> at equilibrium. **(3 marks)** 38 Consider the following equilibrium:

(4 marks)

 $H_{2(g)} + I_{2(g)} \rightleftharpoons 2HI_{(g)} \qquad K_{eq} = 49$ 

A 1.00 L container is initially filled with 0.180 mol HI. Calculate the concentration of HI at equilibrium. (4 marks)

F7 39 Consider the following equilibrium:

$$H_{2(g)} + I_{2(g)} \rightleftharpoons 2HI_{(g)} \qquad K_{eg} = 7.1 \times 10^2$$

At equilibrium, the  $[H_2] = 0.012 \text{ mol/L}$  and [HI] = 0.40 mol/L. What is the equilibrium concentration of  $I_2$ ? (2 marks)

F7 40 Consider the following equilibrium:

$$H_{2(g)} + S_{(s)} \rightleftharpoons H_2 S_{(g)} \qquad K_{eq} = 6.8 \times 10^{-2}$$

A 1.0 L container is initially filled with 0.050 mol  $H_2$  and 0.050 mol S. The container is heated to 90° C and equilibrium is established. What is the equilibrium  $[H_2S]$ ? (3 marks)

F7 41 Consider the following:

 $H_{2(g)} + F_{2(g)} \rightleftharpoons 2HF_{(g)} \qquad K_{eg} = 1.00 \times 10^2$ 

A 1.00 L flask is initially filled with 2.00 mol  $\rm H_2$  and 2.00 mol  $\rm F_2$  .

- Calculate the [H<sub>2</sub>] at equilibrium. (4 marks)
- F7 42 Consider the following equilibrium:

$$2 \operatorname{HI}_{(g)} \rightleftharpoons \operatorname{H}_{2(g)} + \operatorname{I}_{2(g)} \qquad \operatorname{K}_{eq} = 81.0$$

A 1.00 L container is initially filled with 4.00 mol HI. Calculate the [HI] at equilibrium. (4marks)

F7 43 Consider the following equilibrium:

$$N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} \qquad K_{eq} = 626 \text{ at } 200^{\circ}C$$

At equilibrium,  $[N_2]$  is 1.06 mol/L and  $[H_2]$  is 0.456 mol/L. Calculate  $[NH_3]$  in the equilibrium mixture. (2 marks)

F7

F7 44 Consider the data obtained for the following equilibrium:

	$\operatorname{Fe}^{}_{(aq)} + \operatorname{SCN}^{}_{(aq)} \leftarrow \operatorname{FeSCN}^{}_{(aq)}$		
	[Fe <sup>3+</sup> ]	[SCN <sup>-</sup> ]	$\left[\text{FeSCN}^{2+}\right]$
Experiment 1	$3.91 \times 10^{-2}$	$8.02 \times 10^{-5}$	$9.22 \times 10^{-4}$
Experiment 2	$6.27 \times 10^{-3}$	$3.65 \times 10^{-4}$	?

CONT

- 3+

Calculate the  $[FeSCN^{2+}]$  in experiment #2.

(3 marks)

### F8 45 Consider the following equilibrium system:

$$C_{(s)} + 2H_{2(g)} \rightleftharpoons CH_{4(g)} \qquad K_{eq} = 8.1 \times 10^8 \text{ at } 25^{\circ}C$$

A student places 4.5 mol of carbon,  $3.6 \times 10^{-3}$  mol of hydrogen and 5.1 mol of methane in a 1.0 L flask. The student predicts that the [CH<sub>4</sub>] increases as equilibrium is established. Do you agree? Explain your answer using appropriate calculations. **(3 marks)** 

F8 46 Consider the following equilibrium system:

$$C_{(s)} + H_2 O_{(g)} \rightleftharpoons CO_{(g)} + H_{2(g)} \qquad K_{eq} = 0.80$$

In an experiment, a student places 0.10 mol of C, 0.15 mol of  $H_2O$ , 0.25 mol of CO, and 0.20 mol of  $H_2$  into a 1.0 L flask. The student predicts that the [CO] will decrease as equilibrium becomes established. (3 marks)

a) Would you agree or disagree with the student?

b) Justify your answer, including appropriate calculations.

E5 47 The production of ammonia by the Haber process involves the following equilibrium:

$$N_{2(g)} + 3H_{2(g)} \rightleftharpoons 2NH_{3(g)} + heat$$

The table below indicates the percentage of ammonia in equilibrium mixtures at various temperatures.

Temperature °C	Percentage of Ammonia in Equilibrium		
200	98		
350	80		
500	51		

a) Explain why the lower temperature results in a higher percentage of ammonia in the equilibrium mixture. (1 mark)

b) Explain why a temperature of 500°C is used in the Haber process rather than a lower temperature. (1 mark)

E	QU	ILIBRIUM WRITTEN SOLUTIONS (FROM PROVINCIAL KEYS)				
D3	1	[NOCI] decreases as it approaches equilibrium. [Cl <sub>2</sub> ] increases as it approaches equilibrium. <b>1 mark</b>				
D4	2	Closed container.				
	Constant temperature.					
		Reversible reaction.				
		Both reactants and products present.				
		No changes in macroscopic properties.				
		Rate of forward reaction equals rate of reverse reaction.				
		Responds to imposed stresses.     any four for1/2 mark each				
D4	3	The rates of the forward and reverse reactions.				
D5	4	a) Both the forward and reversed reactions continue to occur. <b>1mark</b>				
		b) A chemical system at equilibrium is recognized by its constant macroscopic properties.1 mark				
D7	5	a) Entropy is decreasing. Five particles of gas (reactants) have more entropy than four particles of gas				
		(products).1 mark				
		b) Enthalpy is decreasing. The reaction is exothermic, so the enthalpy of the				
		products is less than the enthalpy of the reactants. <b>1 mark</b>				
D9	6	For Example:				
		Enthalpy: is decreasing. 1/2 mark				
		Entropy: is decreasing. 1/2 mark				
		Explanation: Since the system reaches equilibrium, the drive to minimum enthalpy and maximum				
⊏1	7					
	1	when a system at equilibrium is subjected to a stress, processes occur				
		that tend to counteract the stress and re-establish equilibrium. 2 marks				
E1	8	When a system at equilibrium(1/2 mark) is subjected to a stress, (1/2 mark) the system shifts so as to offset				
50		the stress $(1/2 \text{ mark})$ and establish a new equilibrium $(1/2 \text{ mark})$ .				
E2	9	a) The amount of $Cl_2$ will increase because the equilibrium shifts left. <b>1 mark</b>				
ГO	10	b) The amount of Cl <sub>2</sub> will decrease because the equilibrium shifts fight. 1 mark				
ΕZ	10	a) The moles of $CH_3OH$ will not change because the equilibrium shifts right. <b>A mark</b>				
<b>E</b> 2	11	b) The moles of Ch <sub>3</sub> OH will increase because the equilibrium shifts right. Thatk				
LZ		to shift right				
F2	12					
	14	HCl neutralizes $OH^ H^+ + OH^- \rightarrow H_2O$				
		$\langle -2 \text{ marks} \rangle$				
		$\therefore$ [OH <sup>-</sup> ] decreases, therefore equilibrium shifts right (orange).				
F2	13	a) NO $H_0 O_0$ <b>1/2 mark each</b>				
	10	b) $N_0H_4$ <b>1/2 mark</b>				
E2	14	agree with student $\leftarrow 1/2$ mark				
		agree with student ( 1/2 mark				
		cold water bath caused shift in forward direction $\leftarrow 1/2$ mark				
		when temp. is decreased, equil shifts in exo direction <b>I mark</b>				
E2	15	a)				
		The reduced $[\text{Ee}^{3+}]$ causes a shift to the left to offect				
		The reduced [i.e. ] causes a smit to the left to offset $\left\{ \leftarrow 2 \text{ marks} \right\}$				
		the stress.				
		b) The rate of the reverse reaction decreases. <b>1 mark</b>				





F1 21  

$$H_{2(p)} + I_{2(p)} \neq 2H_{1(p)}$$

$$\left[ \begin{array}{c} |I| \\ x \\ |I| \\ -0.080 \\ \hline \\ |E| \\ x - 0.080 \\ x - 0.080 \\ \hline \\ x - 1\frac{1}{2} \text{ marks}$$

$$\left[ \begin{array}{c} I \\ I \\ I \\ I \\ 0.055 \\ \hline \\ 0.050 \\ \hline \\ 0.050 \\ \hline \\ x - 0.015 \\ \hline \\ x - 1\frac{1}{2} \text{ marks}$$

$$\left[ \begin{array}{c} I \\ I \\ 0.055 \\ I \\ 0.050 \\ \hline \\ x - 0.015 \\ \hline \\ x - 0.$$

F5 25  $K_{eq} = \frac{[\text{isobutane}]}{[n - \text{butane}]} \leftarrow \frac{1}{2} \text{ mark}$  $=\frac{3.0}{2.0}$   $\leftarrow$  1 mark  $= 1.5 \qquad \leftarrow \frac{1}{2} \text{ mark}$ F5 26 a) The concentrations become constant (1 mark) b)  $A \rightleftharpoons 2B + C$  (2 marks) (2marks) C)  $K_{eq} = \frac{[B]^{2}[C]}{[A]} = \frac{(0.40)^{2}(0.20)}{0.60} = 0.053$ F5 27 a) Temperature is increased. 1 mark b) NO<sub>2</sub> added. 1 mark c)  $K_{eq} = \frac{[NO_2]^2}{[N_2O_4]} = \frac{(0.60)^2}{(0.20)} = 1.8 \quad \leftarrow 2 \text{ marks}$ F5 28 +  $O_2 \rightleftharpoons 2NO_2$ 2NO  $\frac{0.044 \text{ mol}}{2.00 \text{ L}} \qquad \frac{0.100 \text{ mol}}{2.00 \text{ L}} \qquad \frac{7.88 \text{ mol}}{2.00 \text{ L}} \\ \left. \left. \right\} \leftarrow 1 \text{ mark}$ [E] = 0.022 = 0.0500 = 3.94 $K_{eq} = \frac{\left[NO_{2}\right]^{2}}{\left[NO\right]^{2}\left[O_{2}\right]} = \frac{(3.94)^{2}}{(0.022)^{2}(0.0500)}$  $\leftarrow 1\frac{1}{2}$  marks  $= 6.4 \times 10^5$  $\leftarrow \frac{1}{2}$  mark F6 29  $2 \text{HBr}_{(g)} \rightleftharpoons \text{H}_{2(g)} + \text{Br}_{2(g)}$  $\begin{bmatrix} \mathbf{I} \\ 0.500 & 0 & 0 \\ \hline \mathbf{C} \\ \hline -0.171 & +0.0855 & +0.0855 \\ \hline \mathbf{E} \\ \hline 0.329 & 0.0855 & 0.0855 \\ \end{bmatrix} \leftarrow \mathbf{1} \frac{1}{2} \text{ marks}$  $\mathbf{K}_{eq} = \frac{\left[\mathbf{H}_{2}\right]\left[\mathbf{Br}_{2}\right]}{\left[\mathbf{HBr}\right]^{2}}$  $\leftarrow 1\frac{1}{2}$  marks  $=\frac{(0.0855)(0.0855)}{(0.329)^2}$  $= 6.75 \times 10^{-2}$ 

F6 31

Deduct  $\frac{1}{2}$  mark for incorrect significant figures.

$$\left. \begin{array}{cccc} 2 \operatorname{CH}_{4(g)} &\rightleftharpoons & \operatorname{C}_{2}\operatorname{H}_{2(g)} &+ & 3\operatorname{H}_{2(g)} \\ \\ \left[ \mathbf{I} \right] & 0.180 & 0 & 0 \\ \\ \hline \left[ \mathbf{C} \right] & -0.160 & +0.0800 & +0.240 \\ \hline \left[ \mathbf{E} \right] & 0.020 & 0.0800 & 0.240 \end{array} \right\} \leftarrow 2 \text{ marks}$$

$$K_{eq} = \frac{[C_2H_2][H_2]^3}{[CH_4]^2}$$
  
=  $\frac{(0.0800)(0.240)^3}{(0.020)^2}$   
= 2.8

$$2CO_{(g)} + O_{2(g)} \rightleftharpoons 2CO_{2(g)}$$

$$\begin{bmatrix} \mathbf{I} \\ 0.200 & 0.400 & 0.000 \\ \hline \mathbf{C} \\ -0.160 & -0.080 & +0.160 \\ \hline \mathbf{E} \\ 0.040 & 0.320 & 0.160 \end{bmatrix} \leftarrow \mathbf{1} \frac{1}{2} \text{ marks}$$
$$\mathbf{K}_{eq} = \frac{\left[ CO_2 \right]^2}{\left[ CO \right]^2 \left[ O_2 \right]} = \frac{\left( 0.160 \right)^2}{\left( 0.040 \right)^2 \left( 0.320 \right)} \\ = 5.0 \times 10^1 \end{bmatrix} \leftarrow \mathbf{1} \frac{1}{2} \text{ marks}$$

 $= 8.44 \times 10^{2}$ 

$$F6 = 36 \qquad [Fe^{3x}] = \frac{50.0 \text{ mL}}{80.0 \text{ mL}} \times 0.10 \text{ M} = 0.0625 \text{ M}} \\ [SCN^{-}] = \frac{30.0 \text{ mL}}{80.0 \text{ mL}} \times 0.20 \text{ M} = 0.0750 \text{ M}} \qquad \left| \leftarrow 1 \text{ mark} \right| \\ [I] = \frac{Fe^{3x}}{0.0625} + SCN^{-2x} \text{ FeSCN}^{2x}} \\ [I] = \frac{10025}{0.0250} - 0.0500} - 0.0500} \\ [C] = -0.0500 - 0.0500} \\ [E] = 0.0125 - 0.0250} \\ = \frac{10025}{0.0125} - 0.0250} \\ = \frac{1}{12} \text{ marks}} \\ F7 = \frac{11}{2} \text{ marks}} \\ F7 = \frac{11}{2} \text{ marks}} \\ [K_{eq}] = \frac{[H1]^2}{[H_1][L_2]} = 1.2 \times 10^{-2}} \\ = \frac{0.0500}{(0.0125)(0.0250)} \\ = 1.6 \times 10^{2}} \\ F7 = \frac{11}{2} \text{ marks}} \\ K_{eq}] = \frac{[H1]^2}{[H_1][L_2]} = 1.2 \times 10^{-2}} \\ = \frac{(0.050 - 2x)^2}{x^2} = \sqrt{1.2 \times 10^{-2}} \\ = \sqrt{\frac{(0.050 - 2x)^2}{x^2}} = \sqrt{1.2 \times 10^{-2}} \\ = \sqrt{\frac{(0.050 - 2x)^2}{x^2}} = \sqrt{1.2 \times 10^{-2}} \\ = \sqrt{\frac{(0.050 - 2x)^2}{x^2}} = \sqrt{1.2 \times 10^{-2}} \\ K_{eq}] = \frac{[H_2] = 0.024 \text{ M}} \\ \leftarrow \frac{1}{2} \text{ mark} \\ F7 = \frac{1}{38} \\ \frac{H_{2(g)} + 1_{2(g)} \in 2 2 \text{ H}_{1(g)}}{[H] = 0 = 0 \text{ 0.180}} \\ [C] + x + x - -2x \\ [E] = x - x - 0.80 - 2x \\ \end{bmatrix} \leftarrow \frac{1}{2} \text{ mark} \\ \frac{1}{4} \text{ mark} \\ \frac{1}$$

$$K_{eq} = \frac{[HI]^{2}}{[H_{2}][I_{2}]}$$

$$= \frac{(0.180 - 2x)^{2}}{(x)^{2}} = 49$$

$$x = 0.020$$

$$[HI] = 0.180 - 2x$$

$$= 0.140 \text{ mol/L}$$

$$\left\{ \leftarrow 1 \text{ mark} \right\}$$

F7 39  

$$K_{eq} = \frac{[HI]^{2}}{[H_{2}][I_{2}]}$$

$$\therefore [I_{2}] = \frac{[HI]^{2}}{[H_{2}]K_{eq}}$$

$$= \frac{(0.40)^{2}}{(0.012)(7.1 \times 10^{2})}$$

$$= 0.019 \text{ mol/L}$$

$$\leftarrow 2 \text{ marks}$$

 $K_{eq} = \frac{[HF]^2}{[H_2][F_2]}$  $1.00 \times 10^2 = \frac{(2x)^2}{(2.00 - x)^2}$ 

x = 1.67

 $[H_2] = 2.00 - x = 2.00 - 1.67 = 0.33 \text{ mol/L} \leftarrow 1 \text{ mark}$ 

NOTE:	$\left(\frac{1}{2}\right)$	mark)	is deducted	for incorrect	significant	figures.
-------	----------------------------	-------	-------------	---------------	-------------	----------

 $\left. \leftarrow 1\frac{1}{2} \text{ marks} \right.$ 





$$\mathbf{K}_{eq} = \frac{\left[\mathbf{NH}_{3}\right]^{2}}{\left[\mathbf{N}_{2}\right]\left[\mathbf{H}_{2}\right]^{3}} \qquad \leftarrow \frac{1}{2} \text{ mark}$$

$$626 = \frac{\left[\mathrm{NH}_3\right]^2}{(1.06)(0.456)^3} \quad \leftarrow \frac{1}{2} \text{ mark}$$

$$[NH_3] = 7.93$$

 $\leftarrow 1 \text{ mark}$ 

$$K_{eq} = \frac{\left[\text{FeSCN}^{2+}\right]}{\left[\text{Fe}^{3+}\right]\left[\text{SCN}^{-}\right]} = \frac{9.22 \times 10^{-4}}{(3.91 \times 10^{-2})(8.02 \times 10^{-5})} = 2.94 \times 10^{2} \qquad \left\{ \leftarrow 1\frac{1}{2} \text{ mark} \right\}$$

$$2.94 \times 10^{2} = \frac{x}{(6.27 \times 10^{-3})(3.65 \times 10^{-4})}$$
  
[FeSCN<sup>2+</sup>] = x = 6.73 × 10<sup>-4</sup> M   
$$\leftarrow 1\frac{1}{2} \text{ mark}$$

(Deduct  $\frac{1}{2}$  mark for incorrect significant figures.)



### Note: solutions to questions 46,47 intentionally left off the key...hmmmm?